

# Partial Motzkin paths with air pockets of the first kind avoiding peaks, valleys or double rises

Jean-Luc Baril<sup>1</sup> and José L. Ramírez<sup>2</sup>

<sup>1</sup>LIB, Université de Bourgogne  
B.P. 47 870, 21078 Dijon Cedex France

E-mail: `barjl@u-bourgogne.fr`

<sup>2</sup>Departamento de Matemáticas, Universidad Nacional de Colombia  
Bogotá, Colombia

E-mail: `jlr Ramirezr@unal.edu.co`

July 18, 2024

## Abstract

Motzkin paths with air pockets (MAP) of the first kind are defined as a generalization of Dyck paths with air pockets. They are lattice paths in  $\mathbb{N}^2$  starting at the origin made of steps  $U = (1, 1)$ ,  $D_i = (1, -i)$ ,  $i \geq 1$  and  $H = (1, 0)$ , where two down-steps cannot be consecutive. We enumerate MAP and their prefixes avoiding peaks (resp. valleys, resp. double rise) according to the length, the type of the last step, and the height of its end-point. We express our results using Riordan arrays. Finally, we provide constructive bijections between these paths and restricted Dyck and Motzkin paths.

## 1 Introduction

In a recent paper [6], the authors introduce, study, and enumerate special classes of lattice paths, called *Dyck paths with air pockets* (DAP for short). Such paths are non empty lattice paths in the first quadrant of  $\mathbb{Z}^2$  starting at the origin, and consisting of up-steps  $U = (1, 1)$  and down-steps  $D_i = (1, -i)$ ,  $i \geq 1$ , where two down-steps cannot be consecutive. These paths can be viewed as ordinary Dyck paths (i.e., paths in  $\mathbb{N}^2$  starting at the origin, ending on the  $x$ -axis and consisting of  $U$  and  $D_1(= D)$ ), where each maximal run of down-steps is condensed into one large down-step. As mentioned in [6], they also correspond to a stack evolution with (partial) reset operations that cannot be consecutive (see for instance [13]). The authors enumerate these paths with respect to the length, the type (up or down) of the

last step and the height of the end-point. Whenever the last point is on the  $x$ -axis, they prove that the DAP of length  $n$  are in one-to-one correspondence with the peak-less Motzkin paths of length  $n - 1$ . They also investigate the popularity of many patterns in these paths and they give asymptotic approximations. In a second work [7], the authors make a study for a generalization of these paths by allowing them to go below the  $x$ -axis. They call these paths Grand Dyck paths with air pockets (GDAP), and they also yield enumerative results for these paths according to the length and several restrictions on the height. In a third paper, Baril and Barry [8] study two generalizations of DAP by allowing some horizontal steps  $H = (1, 0)$  with some conditions. They call them *Motzkin paths with air pockets of the first and second kind*. For other generalizations of Motzkin paths, we refer to [17, 15] for example.

In this paper we study Motzkin paths with air pockets of the first kind, which are defined as Motzkin paths [9] (lattice paths in  $\mathbb{N}^2$  starting at the origin and made of  $U$ ,  $D$ , and  $H$ ), where each maximal run of down-steps is condensed into one large down-step. More precisely, we consider lattice paths in  $\mathbb{N}^2$  starting at the origin, consisting of steps  $U$ ,  $H$ , and  $D_i$ ,  $i \geq 1$ , where *two down-steps cannot be consecutive*. We denote by  $\mathcal{D}$  (resp.  $\mathcal{M}$ , resp.  $\mathcal{MP}$ ) the set of Dyck paths (resp. Motzkin paths, resp. Motzkin paths with air pockets of the first kind). Moreover, we denote by  $\mathcal{PMP}$  the set of partial Motzkin paths with air pockets (PMAP for short). The MAP of the first kind are enumerated by the sequence [A114465](#). This sequence also counts the Dyck paths having no ascents of length 2 that start at an odd level.

Throughout the paper, we will use the following notations. For  $k \geq 0$ , we consider the generating function  $f_k = f_k(z)$  (resp.  $g_k = g_k(z)$ , resp.  $h_k = h_k(z)$ ), where the coefficient of  $z^n$  in the series expansion is the number of partial Motzkin paths with air pockets of length  $n$  ending at height  $k$  with an up-step, (resp. with a down-step, resp. with a horizontal step  $H$ ). We introduce the bivariate generating functions

$$F(u, z) = \sum_{k \geq 0} u^k f_k(z), \quad G(u, z) = \sum_{k \geq 0} u^k g_k(z), \quad \text{and} \quad H(u, z) = \sum_{k \geq 0} u^k h_k(z).$$

For short, we also use the notation  $F(u)$ ,  $G(u)$ , and  $H(u)$  for these functions.

A *Riordan array* is an infinite lower triangular matrix whose  $k$ -th column has generating function  $g(z)f(z)^k$  for all  $k \geq 0$ , for some formal power series  $g(z)$  and  $f(z)$ , with  $g(0) \neq 0$ ,  $f(0) = 0$ , and  $f'(0) \neq 0$ . Such a Riordan array is denoted by  $(g(z), f(z))$ . We refer to [25, 26] for more details on Riordan arrays. Several authors have used Riordan arrays to study lattice paths; see for example [19, 22, 23, 29, 30, 31].

The outline of this paper is the following. We present enumerative results for partial Motzkin paths with air pockets of the first kind avoiding peaks (resp. avoiding valleys, resp. avoiding double rises), knowing that a *peak* is an occurrence  $UD_i$  for some  $i \geq 1$ , a *valley* is an occurrence  $D_iU$  for some  $i \geq 1$ , and a *double rise* is an occurrence  $UU$ . For each avoidance, we provide bivariate generating functions that count the PMAP with respect to the length, the type of the last step (up, down or horizontal step) and the height of the end-point. All these results are obtained algebraically by using the famous kernel method for solving several systems of functional equations [1, 20]. We express our results using Riordan

arrays and we deduce closed forms for PMAP of length  $n$  ending at height  $k$ . Finally, we provide constructive bijections between these paths and some restricted Dyck and Motzkin paths.

## 2 Partial peak-less Motzkin paths with air pockets

In this section, we study partial Motzkin paths with air pockets of the first kind avoiding occurrences of  $UD_i$  for all  $i \geq 1$ .

### 2.1 Enumerative results

Let  $P$  be a length  $n$  PMAP ending at height  $k \geq 0$  and avoiding the occurrences of  $UD_i$  for  $i \geq 1$ . If the last step of  $P$  is  $U$ , then  $k \geq 1$  and we have  $P = QU$ , where  $Q$  is a length  $(n-1)$  MAP ending at height  $k-1$  and avoiding the peaks ( $Q$  can be the empty path). So, we obtain the first relation  $f_k = zf_{k-1} + zg_{k-1} + zh_{k-1}$  for  $k \geq 1$ , anchored with  $f_0 = 1$  by considering the empty path. If the last step of  $P$  is a down-step  $D_i$ ,  $i \geq 1$ , then we have  $P = QD_i$ , where  $Q$  is a length  $(n-1)$  PMAP ending at height  $\ell \geq k+1$  with no up- and down-steps at its end, and with no peaks. So, we obtain the second relation  $g_k = z \sum_{\ell \geq k+1} h_\ell$ .

If the last step of  $P$  is a horizontal step  $H$ , then we have  $P = QH$ , where  $Q$  is a length  $(n-1)$  PMAP ending at height  $k$  with no peaks, which implies that  $h_k = z(f_k + g_k + h_k)$  for  $k \geq 0$ .

Therefore, we have to solve the following system of equations:

$$\begin{cases} f_0 = 1, \text{ and } f_k = zf_{k-1} + zg_{k-1} + zh_{k-1}, & k \geq 1, \\ g_k = z \sum_{\ell \geq k+1} h_\ell, & k \geq 0, \\ h_k = zf_k + zg_k + zh_k, & k \geq 0. \end{cases} \quad (1)$$

Multiplying by  $u^k$  the recursions in (1) and summing over  $k$ , we have:

$$\begin{aligned} F(u) &= 1 + z \sum_{k \geq 1} u^k f_{k-1} + z \sum_{k \geq 1} u^k g_{k-1} + z \sum_{k \geq 1} u^k h_{k-1} \\ &= 1 + zuF(u) + zuG(u) + zuH(u), \\ G(u) &= z \sum_{k \geq 0} u^k \left( \sum_{\ell \geq k+1} h_\ell \right) = z \sum_{k \geq 1} h_k (1 + u + \dots + u^{k-1}) \\ &= z \sum_{k \geq 1} \frac{u^k - 1}{u - 1} h_k = \frac{z}{u - 1} (H(u) - H(1)), \\ H(u) &= zF(u) + zG(u) + zH(u). \end{aligned}$$

Notice that we have  $F(1) - H(1) = 1$  by considering the difference of the first and third equations. Now, setting  $h1 := H(1)$  and solving these functional equations, we obtain

$$F(u) = \frac{h1 uz^2 + zu + z^2 - u - z + 1}{u^2z + z^2 - u - z + 1},$$

$$G(u) = -\frac{z(h1 uz + zh1 - h1 + z)}{u^2z + z^2 - u - z + 1}, \quad H(u) = \frac{z(zh1 - u + 1)}{u^2z + z^2 - u - z + 1}.$$

In order to compute  $h1$ , we use the kernel method (see [2, 20]) on  $H(u)$ . We can write the denominator (which is a polynomial in  $u$  of degree 2), as  $z(u-r)(u-s)$  with

$$r = \frac{1 + \sqrt{-4z^3 + 4z^2 - 4z + 1}}{2z} \quad \text{and} \quad s = \frac{1 - \sqrt{-4z^3 + 4z^2 - 4z + 1}}{2z}.$$

Plugging  $u = s$  (which has a Taylor expansion at  $z = 0$ ) in  $H(u)z(u-r)(u-s)$ , we obtain the equation  $zh1 - s + 1 = 0$ , which implies that

$$h1 = \frac{s-1}{z}.$$

Finally, after simplifying by the factor  $(u-s)$  in the numerators and denominators, we obtain

$$F(u) = \frac{r}{r-u}, \quad G(u) = \frac{s-1}{r-u}, \quad \text{and} \quad H(u) = \frac{1}{r-u},$$

which induces that

$$f_k = [u^k]F(u) = \frac{1}{r^k}, \quad g_k = [u^k]G(u) = \frac{s-1}{r^{k+1}}, \quad \text{and} \quad h_k = [u^k]H(u) = \frac{1}{r^{k+1}}.$$

It is worth noticing that the system of equations for  $F(u)$ ,  $G(u)$ , and  $H(u)$  can be solved directly using the algorithm presented in [12].

**Theorem 1** *The bivariate generating function for the total number of peak-less PMAP with respect to the length and the height of the end-point is given by*

$$Total(z, u) = \frac{1}{z(r-u)},$$

and we have

$$[u^k]Total(z, u) = \frac{1}{zr^{k+1}}.$$

Finally, setting  $t(n, k) = [z^n][u^k]Total(z, u)$ , we have for  $n \geq 2$  and  $k \geq 1$ ,

$$t(n, k) = t(n, k-1) + t(n-1, k) - t(n-1, k-2) - t(n-2, k),$$

and setting  $t_n := t(n, 0)$ , then we have

$$t_n = t_{n-1} + \sum_{k=0}^{n-3} t_k t_{n-k-3} + \sum_{k=2}^{n-1} (t_k - t_{k-1}) t_{n-k-1}.$$

*Proof.* The first three equalities are immediately deduced from the previous results. Now, let us prove the last equality. Any length  $n$  peak-less MAP is of the form (i)  $HP$  where  $P$  is a MAP of length  $n - 1$ , or (ii)  $UQHDR$ , where  $Q, R$  are some MAP such that the length of  $Q$  lies into  $[0, n - 3]$ , or (iii)  $P^\sharp Q$ , where  $P^\sharp = UP'D_i$ ,  $i \geq 2$ , and  $P'D_{i-1}$  is a MAP of length lying into  $[2, n - 1]$ . The number of  $P'D_{i-1}$  of a given length  $k$  is the total number of peak-less MAP of length  $k$  minus the number of peak-less MAP of length  $k$  and ending with  $H$ . Taking into account all these cases, we obtain the result.  $\square$

**Corollary 1** *The generating function that counts all peak-less PMAP with respect to the length is given by*

$$Total(z, 1) = \frac{1}{z(r-1)}.$$

The first few terms of the series expansion of  $Total(z, 1)$  are

$$1 + 2z + 4z^2 + 9z^3 + 22z^4 + 56z^5 + 146z^6 + 388z^7 + 1048z^8 + 2869z^9 + O(z^{10}),$$

which corresponds to the sequence [A152225](#) in [27] counting Dyck paths of semilength  $n + 1$  with no peaks of height  $0 \pmod{3}$  and no valleys of height  $2 \pmod{3}$ ; see [14].

**Corollary 2** *The generating function that counts the peak-less MAP with respect to the length is given by*

$$Total(z, 0) = \frac{1}{zr}.$$

The first few terms of the series expansion of  $Total(z, 0)$  are

$$1 + z + z^2 + 2z^3 + 5z^4 + 12z^5 + 29z^6 + 73z^7 + 190z^8 + 505z^9 + O(z^{10}),$$

which corresponds to the sequence [A152171](#) in [27] counting Dyck paths of length  $2n$  with no peaks of height  $2 \pmod{3}$  and no valleys of height  $1 \pmod{3}$ . In Section 2.2, we will exhibit a constructive bijection between these two classes of paths.

Let  $\mathcal{T}$  be the infinite matrix  $\mathcal{T} := [t(n, k)]_{n, k \geq 0}$ , where  $t(n, k) = [z^n][u^k] Total(z, u)$ . The first few rows of the matrix  $\mathcal{T}$  are

$$\mathcal{T} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 3 & 3 & 1 & 0 & 0 & 0 & 0 & 0 \\ 5 & 6 & 6 & 4 & 1 & 0 & 0 & 0 & 0 \\ 12 & 15 & 13 & 10 & 5 & 1 & 0 & 0 & 0 \\ 29 & 38 & 33 & 24 & 15 & 6 & 1 & 0 & 0 \\ 73 & 96 & 87 & 63 & 40 & 21 & 7 & 1 & 0 \\ 190 & 248 & 229 & 172 & 110 & 62 & 28 & 8 & 1 \end{pmatrix}.$$

In Figure 1 we show the peak-less PMAP counted by  $t(5, 1) = 15$ .




Figure 1: Peak-less PMAP of length 5 ending at height 1.

**Corollary 3** *The matrix  $\mathcal{T} = [t(n, k)]_{n, k \geq 0}$  is a Riordan array defined by*

$$(C(z(1-z+z^2)), zC(z(1-z+z^2))),$$

where  $C(z) = \frac{1-\sqrt{1-4z}}{2z}$  is the generating function of the Catalan numbers  $c_n = \frac{1}{n+1} \binom{2n}{n}$ .

*Proof.* Indeed, we directly deduce the result from the following.

$$[u^k] \text{Total}(z, u) = \frac{1}{zr^{k+1}} = \frac{1}{zr} \cdot \frac{1}{r^k} = C(z(1-z+z^2)) \cdot (zC(z(1-z+z^2)))^k.$$

□

**Corollary 4** *We have*

$$t(n, k) = \sum_{j=0}^{n-k} \frac{k+1}{2(n-j)-k+1} \binom{2(n-j)-k+1}{n-k-j} a(n-k-j, j),$$

where  $a(n, k) = (-1)^k \sum_{i=0}^n \binom{n}{i} \binom{n-i}{k-2i}$ .

*Proof.* From the Lagrange Inversion Formula (cf. [16, 28]), we know that

$$C(z)^k = \sum_{n \geq 0} \frac{k}{2n+k} \binom{2n+k}{n} z^n. \quad (2)$$

From definition of Riordan arrays and Eq. (2) we have

$$\begin{aligned}
t(n, k) &= [z^{n-k}] \left( C(z(1-z+z^2)) \right)^{k+1} \\
&= [z^{n-k}] \sum_{\ell=0}^{\infty} \frac{k+1}{2\ell+k+1} \binom{2\ell+k+1}{\ell} z^\ell (1-z+z^2)^\ell \\
&= [z^{n-k}] \sum_{\ell=0}^{\infty} \frac{k+1}{2\ell+k+1} \binom{2\ell+k+1}{\ell} z^\ell \sum_{j=0}^{2\ell} a(\ell, j) z^j \\
&= [z^{n-k}] \sum_{j=0}^{\infty} \sum_{\ell=0}^{\infty} \frac{k+1}{2(\ell+\lfloor j/2 \rfloor)+k+1} \binom{2(\ell+\lfloor j/2 \rfloor)+k+1}{\ell+\lfloor j/2 \rfloor} a(\ell+\lfloor j/2 \rfloor, j) z^{j+\ell+\lfloor j/2 \rfloor}.
\end{aligned}$$

If we take  $s = j + \ell + \lfloor j/2 \rfloor$ , then

$$\begin{aligned}
t(n, k) &= [z^{n-k}] \sum_{j=0}^{\infty} \sum_{s=j}^{\infty} \frac{k+1}{2(s-j)+k+1} \binom{2(s-j)+k+1}{s-j} a(s-j, j) z^s \\
&= \sum_{j=0}^{n-k} \frac{k+1}{2(n-j)-k+1} \binom{2(n-j)-k+1}{n-k-j} a(n-k-j, j).
\end{aligned}$$

□

In [24], Rogers gave an equivalent characterization of the Riordan arrays. That is, every element not belonging to row 0 or column 0 in a Riordan array can be expressed as a fixed linear combination of the elements in the preceding row. The *A-sequence* is defined to be the sequence coefficients of this linear combination. Analogously, Merlini et al. [18] introduced the *Z-sequence*, that characterizes the elements in column 0, except for the top one.

An infinite lower triangular matrix  $[d_{n,k}]_{n,k \geq 0}$  is a Riordan array if and only if  $d_{0,0} \neq 0$  and there exist two sequences  $(a_0, a_1, a_2, \dots)$ , with  $a_0 \neq 0$ , and  $(z_0, z_1, z_2, \dots)$  (called the *A-sequence* and the *Z-sequence*, respectively), such that

$$d_{n+1,k+1} = a_0 d_{n,k} + a_1 d_{n,k+1} + a_2 d_{n,k+2} + \dots \quad \text{for } n, k \geq 0, \quad (3)$$

$$d_{n+1,0} = z_0 d_{n,0} + z_1 d_{n,1} + z_2 d_{n,2} + \dots \quad \text{for } n \geq 0. \quad (4)$$

The product of two Riordan arrays  $(g(z), f(z))$  and  $(h(z), l(z))$  is defined by

$$(g(z), f(z)) * (h(z), l(z)) = (g(z)h(f(z)), l(f(z))). \quad (5)$$

Under this operation, the set of all Riordan arrays is a group [25]. The identity element is  $I = (1, z)$  and the inverse of  $(g(z), f(z))$  is given by

$$(g(z), f(z))^{-1} = (1/(g \circ f^{<-1>})(z), f^{<-1>}(z)), \quad (6)$$

where  $f^{<-1>}(z)$  denotes the compositional inverse of  $f(z)$ .

The generating functions for the  $A$ -sequence and  $Z$ -sequence of the Riordan array  $\mathcal{F} = (g(z), f(z))$ , with inverse  $\mathcal{F}^{-1} = (d(z), h(z))$ , are given by ([18, 11])

$$A(z) = \frac{z}{h(z)} \quad \text{and} \quad Z(z) = \frac{1}{h(z)} (1 - d_{0,0}d(z)),$$

respectively.

From the definition of the  $A$ -sequence and  $Z$ -sequence for the Riordan arrays we can give an additional recurrence relation for the sequence  $t(n, k)$ .

**Corollary 5** *We have*

$$t(n+1, k+1) = \sum_{j \geq 0} a(j)t(n, k+j),$$

where  $a(n) = (-1)^{n+1} \sum_{k=1}^n \sum_{j=0}^k \frac{1}{k} \binom{j}{n-k-j} \binom{k}{j} \binom{n-k-2}{k-1}$  for  $n \geq 1$  and  $a(0) = 1$ . Moreover,

$$t_{n+1} = \sum_{j \geq 0} a(j+1)t(n, j).$$

*Proof.* By Equation (6), the inverse of the matrix  $\mathcal{T} = [t(n, k)]_{n, k \geq 0}$  is given by  $\mathcal{T}^{-1} = (g_2(z), zg_2(z))$ , where

$$g_2(z) = \frac{-1 + z^2 + \sqrt{1 - 2z^2 + 4z^3 - 3z^4}}{2z^3}.$$

Therefore, the  $A$ -sequence and  $Z$ -sequence of the Riordan array  $\mathcal{T}$  have generating functions

$$A(z) = \sum_{n \geq 0} a(n)z^n = \frac{2z^3}{-1 + z^2 + \sqrt{1 - 2z^2 + 4z^3 - 3z^4}} \quad \text{and} \quad Z(z) = \frac{A(z) - 1}{z}.$$

The generating function  $A(z)$  corresponds with the sequence [A247162](#), where the explicit formula for  $a(n)$  can be found. From (3) we obtain the result.  $\square$

The first few values of the sequence  $a(n)$  for  $n \geq 0$  are

$$1, \quad 1, \quad 0, \quad 1, \quad 0, \quad 1, \quad -1, \quad 2, \quad -3, \quad 6, \quad -10, \dots$$

## 2.2 A bijective approach

Corollary 2 proves that the set of peak-less Motzkin paths with air pockets of length  $n$  (ending on the  $x$ -axis) is equinumerous to the set  $\mathcal{D}_n(2, 1)$  of Dyck paths of length  $2n$  with no peak at height 2 (mod 3) and no valley at height 1 (mod 3).

Any non-empty peak-less Motzkin path with air pockets is either of the form (1)  $H\alpha$  or (2)  $U\alpha_1U\alpha_2 \cdots U\alpha_kHD_k\beta$ , where  $k \geq 1$  and  $\alpha, \alpha_1, \dots, \alpha_k, \beta$  are possibly empty peak-less MAP. We refer to the left part of Figure 2 for an illustration of this form.

Remember that  $\mathcal{MP}$  denotes the set of Motzkin paths with air pockets of the first kind.



**Definition 1** We recursively define the map  $\psi$  from  $\mathcal{MP}$  to  $\cup_{n \geq 0} \mathcal{D}_n(2, 1)$  as follows. For  $\alpha \in \mathcal{MP}$ , we set:

$$\psi(P) = \begin{cases} \epsilon & \text{if } P = \epsilon & (i) \\ UD\psi(\alpha) & \text{if } P = H\alpha \text{ with } \alpha \in \mathcal{MP} & (ii) \\ U^3\psi(\alpha_1)DU\psi(\alpha_2)DU \dots DU\psi(\alpha_k)D^3\psi(\beta) & \text{if } P = U\alpha_1U\alpha_2 \dots U\alpha_kHD_k\beta \text{ with} & (iii) \\ & k \geq 1 \text{ and } \alpha_1, \dots, \alpha_k, \beta \in \mathcal{MP}. \end{cases}$$

We refer to Figure 3 for an illustration of the third case of the definition of  $\psi$ .




Figure 2: Illustration of the map  $\psi$  for the more general case (iii) in Definition 1.

Due to the recursive definition, the image of peak-less MAP of length  $n$  under  $\psi$  is a Dyck path of length  $2n$ . Moreover it is clear that the obtained path is a Dyck path in  $\cup_{n \geq 0} \mathcal{D}_n(2, 1)$ . For instance (see Figure 3 for an illustration of this example).

$$\begin{aligned} \psi(UUHDHUHUHD_3HH) &= \psi(U \cdot \overbrace{UHDH}^{\alpha_1} \cdot U \cdot \overbrace{H}^{\alpha_2} \cdot U \cdot \overbrace{\epsilon}^{\alpha_3} \cdot HD_3 \cdot \overbrace{HH}^{\beta}) \\ &= U^3\psi(UHDH) \cdot DU \cdot \psi(H) \cdot DU \cdot \psi(\epsilon) \cdot D^3 \cdot \psi(HH) \\ &= U^3 \cdot (U^3D^3\psi(H)) \cdot DU \cdot UD \cdot DU \cdot D^3 \cdot UDUD \\ &= U^6D^3UD^2U^2D^2UD^2DUDUD. \end{aligned}$$




Figure 3:  $\psi(UUHDHUHUHD_3HH) = UUUUUDDDUDDUDDUDDDUDDUDD$ .

**Theorem 2** For all  $n \geq 0$ , the map  $\psi$  induces a bijection between  $\mathcal{MP}_n$  and  $\mathcal{D}_n(2, 1)$ .

*Proof.* Since  $\mathcal{MP}_n$  and  $\mathcal{D}_n(2, 1)$  have the same cardinality (due to Corollary 2 and A152171 in [27]), it suffices to prove that for  $P, Q \in \mathcal{MP}$ ,  $P \neq Q$  implies  $\psi(P) \neq \psi(Q)$ . A simple induction on  $n$  allows to obtain the result.  $\square$

### 3 Partial valley-less Motzkin paths with air pockets

Using the same arguments we used for the system of the previous section, we study partial Motzkin paths with air pockets of the first kind avoiding occurrences of  $D_iU$  for all  $i \geq 1$ .

#### 3.1 Enumerative results

In the same way as we done in Section 2.1, we have to solve the following system of equations:

$$\begin{cases} f_0 = 1, \text{ and } f_k = zf_{k-1} + zh_{k-1}, & k \geq 1, \\ g_k = z \sum_{\ell \geq k+1} f_\ell + z \sum_{\ell \geq k+1} h_\ell, & k \geq 0, \\ h_k = zf_k + zg_k + zh_k, & k \geq 0. \end{cases} \quad (7)$$

Multiplying by  $u^k$  the recursions in (7) and summing over  $k$ , we have:

$$\begin{aligned} F(u) &= 1 + zuF(u) + zuH(u), \\ G(u) &= \frac{z}{u-1}(F(u) - F(1) + H(u) - H(1)), \\ H(u) &= zF(u) + zG(u) + zH(u). \end{aligned}$$

This system of equations for  $F(u)$ ,  $G(u)$ , and  $H(u)$  can be solved directly using the algorithm presented in [12]. However we provide the main steps for solving it.

Notice that we have  $(1-z)F(1) = 1 + zH(1)$  by considering the first equation. Now, setting  $f1 := F(1)$  and solving these functional equations, we obtain

$$\begin{aligned} F(u) &= \frac{f1 \, uz^2 - uz^2 + zu + z^2 - u - z + 1}{u^2z + z^2 - u - z + 1}, \\ G(u) &= -\frac{f1 \, uz + f1 \, z - zu - f1 + 1}{u^2z + z^2 - u - z + 1}, \quad H(u) = -\frac{z(f1 \, uz - zu - f1 + u + z)}{u^2z + z^2 - u - z + 1}. \end{aligned}$$

In order to compute  $f1$ , we use the kernel method (see [2, 20]) on  $F(u)$ . We can write the denominator (which is a polynomial in  $u$  of degree 2), as  $z(u-r)(u-s)$  with

$$r = \frac{1 + \sqrt{-4z^3 + 4z^2 - 4z + 1}}{2z} \quad \text{and} \quad s = \frac{1 - \sqrt{-4z^3 + 4z^2 - 4z + 1}}{2z}.$$

Plugging  $u = s$  (which has a Taylor expansion at  $z = 0$ ) in  $F(u)z(u-r)(u-s)$ , we obtain the equation  $f1 \, sz^2 - sz^2 + zs + z^2 - s - z + 1 = 0$ , which implies that

$$f1 = 1 + \frac{s-1}{z}.$$

Finally, after simplifying by the factor  $(u-s)$  in the numerators and denominators, we obtain

$$F(u) = \frac{r}{r-u}, \quad G(u) = \frac{s-1}{z(r-u)}, \quad \text{and} \quad H(u) = \frac{s}{r-u},$$

which implies that

$$f_k = [u^k]F(u) = \frac{1}{r^k}, \quad g_k = [u^k]G(u) = \frac{s-1}{zr^{k+1}}, \quad \text{and} \quad h_k = [u^k]H(u) = \frac{s}{r^{k+1}}.$$

**Theorem 3** *The bivariate generating function for the total number of partial valley-less MAP with respect to the length and the height of the end-point is given by*

$$Total(z, u) = \frac{s}{z(r-u)},$$

and we have

$$[u^k]Total(z, u) = \frac{s}{zr^{k+1}}.$$

Finally, setting  $t(n, k) = [z^n][u^k]Total(z, u)$ , we have for  $n \geq 2$ ,  $k \geq 1$ ,

$$t(n, k) = t(n, k-1) + t(n-1, k) - t(n-1, k-2) - t(n-2, k),$$

and setting  $t_n := t(n, 0)$  and  $t_{-1} = 1$ , then we have for  $n \geq 2$ ,

$$t_{n-1} = t_{n-2} + \sum_{k=0}^{n-3} t_{k-1}t_{n-k-4} + \sum_{k=2}^{n-1} (t_{k-1} - t_{k-2})t_{n-k-2}.$$

*Proof.* The first three equalities are immediately deduced from the previous results. For the last equality, the term  $t_n$  satisfies the same recurrence relation as in Theorem 1 (modulo shift of  $n$ ) since the two sequences are equal modulo a shift.  $\square$

**Corollary 6** *The generating function that counts the partial valley-less MAP with respect to the length is given by*

$$Total(z, 1) = \frac{s}{z(r-1)}.$$

The first few terms of the series expansion of  $Total(z, 1)$  are

$$1 + 2z + 5z^2 + 13z^3 + 34z^4 + 90z^5 + 242z^6 + 660z^7 + 1821z^8 + 5073z^9 + O(x^{10}),$$

which does not appear in [27].

**Corollary 7** *The generating function that counts the partial valley-less MAP with respect to the length is given by*

$$Total(z, 0) = \frac{s}{zr}.$$

The first few terms of the series expansion of  $Total(z, 0)$  are

$$1 + z + 2z^2 + 5z^3 + 12z^4 + 29z^5 + 73z^6 + 190z^7 + 505z^8 + 1363z^9 + O(z^{10}),$$

which corresponds to a shift of the sequence [A152171](#) in [27] counting Dyck paths of length  $2n$  with no peaks of height  $2 \pmod 3$  and no valleys of height  $1 \pmod 3$  (see also Section 3.1).

Let  $\mathcal{T}$  be the infinite matrix  $\mathcal{T} := [t(n, k)]_{n, k \geq 0}$ . The first few rows of the matrix  $\mathcal{T}$  are

$$\mathcal{T} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 5 & 4 & 3 & 1 & 0 & 0 & 0 & 0 & 0 \\ 12 & 10 & 7 & 4 & 1 & 0 & 0 & 0 & 0 \\ 29 & 26 & 18 & 11 & 5 & 1 & 0 & 0 & 0 \\ 73 & 67 & 49 & 30 & 16 & 6 & 1 & 0 & 0 \\ 190 & 175 & 133 & 85 & 47 & 22 & 7 & 1 & 0 \\ 505 & 467 & 361 & 241 & 139 & 70 & 29 & 8 & 1 \end{pmatrix}.$$

In Figure 4 we show the valley-less PMAP counted by  $t(4, 0) = 12$ .




Figure 4: The 12  $UU$ -less PMAP of length 4 ending at height 0.

**Corollary 8** *The matrix  $\mathcal{T} = [t(n, k)]_{n, k \geq 0}$  is a Riordan array defined by*

$$\left( (z^2 - z + 1)C(z(1 - z + z^2))^2, zC(z(1 - z + z^2)) \right)$$

where  $C(z) = \frac{1-\sqrt{1-4z}}{2z}$  is the generating function of the Catalan numbers  $c_n = \frac{1}{n+1} \binom{2n}{n}$ .

*Proof.* Indeed, we directly deduce the result from the following.

$$[u^k] \text{Total}(z, u) = \frac{s}{zr^{k+1}} = \frac{s}{zr} \cdot \frac{1}{r^k} = (z^2 - z + 1)C(z(1 - z + z^2))^2 \cdot (zC(z(1 - z + z^2)))^k.$$

□

From a similar argument as in Corollary 4 we obtain the following result.

**Corollary 9** *We have*

$$t(n, k) = \sum_{j=0}^{n-k} \frac{k+2}{2(n-j) - k + 2} \binom{2(n-j) - k + 2}{n - k - j} a(n - k - j + 1, j),$$

where  $a(n, k) = (-1)^k \sum_{i=0}^n \binom{n}{i} \binom{n-i}{k-2i}$ .

### 3.2 A bijective approach

Corollary 7 and Corollary 2 prove that the set of valley-less Motzkin paths with air pockets of length  $n - 1$  (ending on the  $x$ -axis) is equinumerous to the set of peak-less Motzkin paths with air pockets of length  $n$ , which is in one-to-one correspondence with the set  $\mathcal{D}_n(2, 1)$  of Dyck paths of length  $2n$  with no peak at height 2 (mod 3) and no valley at height 1 (mod 3) (see a constructive bijection in Section 2.2). Below, we provide a bijection between valley-less MAP of length  $n - 1$  and peak-less MAP of length  $n$ .

Any valley-less Motzkin path with air pockets is either of the form (i)  $\epsilon$ , (ii)  $\alpha H$ , (iii)  $U\alpha D$ , (iv)  $\beta H U \alpha D$ , (v)  $U\gamma D_k$ , or (vi)  $\beta H U \gamma D_k$ , where  $\alpha, \beta$  are valley-less MAP (possibly empty), and  $\gamma D_{k-1}$  is a valley-less MAP. According to all these cases, we define the map  $\phi$ .

**Definition 2** *We recursively define the map  $\phi$  from valley-less MAP of length  $n - 1$  to peak-less MAP of length  $n$ . Let  $P$  be a valley-less MAP, we set:*

$$\phi(P) = \begin{cases} H & \text{if } P = \epsilon, & (i) \\ \phi(\alpha)H & \text{if } P = \alpha H, & (ii) \\ U\phi(\alpha)D & \text{if } P = U\alpha D, & (iii) \\ \phi(\beta)U\phi(\alpha)D & \text{if } P = \beta H U \alpha D, & (iv) \\ \phi(\alpha D_{k-1})^\# & \text{if } P = U\alpha D_k, & (v) \\ \phi(\beta)\phi(\alpha D_{k-1})^\# & \text{if } P = \beta H U \alpha D_k, & (vi) \end{cases}$$

where the  $\#$ -operator maps a peak-less MAP of the form  $\alpha D_{k-1}$  into the peak-less MAP  $(\alpha D_{k-1})^\# = U\alpha D_k$ .

Due to the recursive definition, the image of valley-less MAP of length  $n - 1$  under  $\phi$  is a peak-less MAP of length  $n$ . The recursive definition naturally induces that  $\phi$  is a bijection. Using the bijection  $\psi$  presented above, we can easily obtain a constructive bijection between valley-less MAP of length  $n - 1$  and Dyck paths of length  $2n$  with no peak at height 2 (mod 3) and no valley at height 1 (mod 3).

## 4 Partial $UU$ -less Motzkin paths with air pockets

In this section, we study partial Motzkin paths with air pockets of the first kind avoiding occurrences of  $UU$ .

### 4.1 Enumerative results

In the same way as we done in the previous section, we have to solve the following system of equations:

$$\begin{cases} f_0 = 1, f_1 = z + zg_0 + zh_0, \text{ and } f_k = zg_{k-1} + zh_{k-1}, & k \geq 2, \\ g_k = z \sum_{\ell \geq k+1} f_\ell + z \sum_{\ell \geq k+1} h_\ell, & k \geq 0, \\ h_k = zf_k + zg_k + zh_k, & k \geq 0. \end{cases} \quad (8)$$

Multiplying by  $u^k$  the recursions in (8) and summing over  $k$ , we have:

$$\begin{aligned} F(u) &= 1 + zu + zuG(u) + zuH(u), \\ G(u) &= \frac{z}{u-1}(F(u) - F(1) + H(u) - H(1)), \\ H(u) &= zF(u) + zG(u) + zH(u). \end{aligned}$$

Again, this system of equations for  $F(u)$ ,  $G(u)$ , and  $H(u)$  can be solved directly using the algorithm presented in [12]. However we provide the main steps for solving it.

Notice that we have  $H(1) = (1+z)(F(1) - 1)$  by considering the first and the last equation. Now, setting  $f1 := F(1)$  and solving these functional equations, we obtain

$$\begin{aligned} F(u) &= \frac{f1uz^3 + 2f1uz^2 + u^2z^2 - u^2z - 2uz^2 + 2uz + z^2 - u - z + 1}{u^2z^2 + uz^3 + uz + z^2 - u - z + 1}, \\ G(u) &= -\frac{z(f1uz^3 + 2f1uz^2 - uz^3 + f1z^2 - uz^2 + f1z + uz - z^2 - 2f1 + 2)}{u^2z^2 + uz^3 + uz + z^2 - u - z + 1}, \\ H(u) &= \frac{z(f1uz^3 + 2f1uz^2 - uz^3 + f1z^2 - u^2z - 2uz^2 + 2f1z + uz - z^2 - u - 2z + 1)}{u^2z^2 + uz^3 + uz + z^2 - u - z + 1}. \end{aligned}$$

In order to compute  $f1$ , we use the kernel method (see [2, 20]) on  $F(u)$ . We can write the denominator (which is a polynomial in  $u$  of degree 2), as  $z^2(u-r)(u-s)$ , with

$$\begin{aligned} r &= \frac{-z^3 - z + 1 + \sqrt{z^6 - 2z^4 + 2z^3 - 3z^2 - 2z + 1}}{2z^2} \quad \text{and} \\ s &= \frac{1 - z - z^3 - \sqrt{z^6 - 2z^4 + 2z^3 - 3z^2 - 2z + 1}}{2z^2}. \end{aligned}$$

Plugging  $u = s$  (which has a Taylor expansion at  $z = 0$ ) in  $F(u)z^2(u - r)(u - s)$ , we obtain the equation  $f_1 s z^3 + 2f_1 s z^2 + s^2 z^2 - s^2 z - 2s z^2 + 2s z + z^2 - u - z + 1 = 0$ , which implies that

$$f_1 = 1 + \frac{s - 1}{z(z + 2)}.$$

Finally, after simplifying by the factor  $(u - s)$  in the numerators and denominators, we obtain

$$F(u) = \frac{z - 1}{z} + \frac{r}{z(r - u)}, \quad G(u) = \frac{s + z}{r - u}, \quad \text{and} \quad H(u) = \frac{zr + 1}{z(r - u)} - 1,$$

which implies that  $f_0 = 1$ ,  $g_0 = \frac{s+z}{r}$ ,  $h_0 = \frac{1}{zr}$  and

$$f_k = [u^k]F(u) = \frac{1}{zr^k}, \quad g_k = [u^k]G(u) = \frac{s + z}{r^{k+1}}, \quad \text{and} \quad h_k = [u^k]H(u) = \frac{1 + zr}{zr^{k+1}}.$$

**Theorem 4** *The bivariate generating function for the total number of  $UU$ -less PMAP with respect to the length and the height of the end-point is given by*

$$\text{Total}(z, u) = \frac{1 + uz}{(r - u)z^2},$$

and we have

$$[u^0]\text{Total}(z, u) = \frac{1}{z^2 r},$$

and for  $k \geq 1$

$$[u^k]\text{Total}(z, u) = \frac{rz + 1}{z^2 r^{k+1}}.$$

Finally, setting  $t(n, k) = [z^n][u^k]\text{Total}(z, u)$ , we have for  $n \geq 2$ ,  $k \geq 1$ ,

$$t(n, k) = t(n, k - 1) + t(n - 1, k) - t(n - 1, k - 1) - t(n - 2, k) \\ - t(n - 2, k - 2) - t(n - 3, k - 1).$$

and setting  $t_n := t(n, 0)$ , then we have

$$t_n = t_{n-1} + t_{n-2} + t_{n-3} + \sum_{k=2}^{n-2} t_{k-2} t_{n-k-2} + \sum_{k=3}^{n-1} (t_{k-1} - t_{k-2}) t_{n-k-1}.$$

*Proof.* The first three equalities are immediately deduced from the previous results. Now, let us prove the last equality. Any non-empty length  $n$   $UU$ -less MAP is of the form (i)  $HP$  where  $P$  is a  $UU$ -less MAP of length  $n - 1$ , or (ii)  $UDQ$  where  $Q$  is a MAP of length  $n - 2$  avoiding  $UU$ , or (iii)  $UQDR$ , where  $Q, R$  are some MAP avoiding  $UU$  such that the length of  $Q$  lies into  $[1, n - 2]$  and  $Q$  starts and ends with  $H$ , or (iv)  $PQ$ , where  $P = UP'D_i$ ,  $i \geq 2$ , and  $P'D_{i-1}$  is a MAP of length lying into  $[3, n - 1]$  and starting with  $H$ . The number of  $P'D_{i-1}$  of a given length  $k$  is the total number of  $UU$ -less MAP of length  $k - 1$  minus the total number of  $UU$ -less MAP of length  $k - 2$ . Taking into account all these cases, we obtain the result. □

**Corollary 10** *The generating function that counts the partial  $UU$ -less MAP with respect to the length is given by*

$$Total(z, 1) = \frac{1+z}{(r-1)z^2}.$$

The first few terms of the series expansion of  $Total(z, 1)$  are

$$1 + 2z + 4z^2 + 9z^3 + 21z^4 + 50z^5 + 122z^6 + 302z^7 + 759z^8 + 1928z^9 + O(x^{10}),$$

which does not appear in [27].

**Corollary 11** *The generating function that counts  $UU$ -less MAP with respect to the length is given by*

$$Total(z, 0) = \frac{1}{z^2 r}.$$

The first few terms of the series expansion of  $Total(z, 0)$  are

$$1 + z + 2z^2 + 4z^3 + 9z^4 + 20z^5 + 47z^6 + 112z^7 + 274z^8 + 679z^9 + O(z^{10}),$$

which corresponds to the sequence [A095980](#) in [27] counting Motzkin paths of length  $n$  with no occurrences of  $UHU$ .

Let  $\mathcal{T}$  be the infinite matrix  $\mathcal{T} := [t(n, k)]_{n, k \geq 0}$ . The first few rows of the matrix  $\mathcal{T}$  are

$$\mathcal{T} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 4 & 4 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 9 & 9 & 3 & 0 & 0 & 0 & 0 & 0 & 0 \\ 20 & 21 & 8 & 1 & 0 & 0 & 0 & 0 & 0 \\ 47 & 50 & 21 & 4 & 0 & 0 & 0 & 0 & 0 \\ 112 & 121 & 55 & 13 & 1 & 0 & 0 & 0 & 0 \\ 274 & 298 & 143 & 39 & 5 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

In Figure 5 we show the  $UU$ -less PMAP counted by  $t(4, 0) = 9$ .




Figure 5: The 9  $UU$ -less PMAP of length 4 ending at height 0.



The matrix  $\mathcal{G}$  is not a (proper) Riordan array. For this reason, we consider the matrix  $\mathcal{G} := [g(n, k)]_{n \geq 0, k \geq 0}$ , where

$$g(n, k) = \begin{cases} 1, & \text{if } n = k = 0; \\ t(n + k - 1, k), & \text{if } n \geq 1. \end{cases}$$

The first few rows of the matrix  $\mathcal{G}$  are

$$\mathcal{G} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 4 & 3 & 1 & 0 & 0 & 0 & 0 & 0 \\ 4 & 9 & 8 & 4 & 1 & 0 & 0 & 0 & 0 \\ 9 & 21 & 21 & 13 & 5 & 1 & 0 & 0 & 0 \\ 20 & 50 & 55 & 39 & 19 & 6 & 1 & 0 & 0 \\ 47 & 121 & 143 & 113 & 64 & 26 & 7 & 1 & 0 \\ 112 & 298 & 372 & 319 & 203 & 97 & 34 & 8 & 1 \end{pmatrix}.$$

**Corollary 12** *The matrix  $\mathcal{G} = [g(n, k)]_{n, k \geq 0}$  is the Riordan array defined by  $(t(z), t(z) - 1)$ , where*

$$t(z) = \frac{1 + z(1 - z)^2 - \sqrt{(1 - 3z + z^3)(1 + z + z^3)}}{2z(1 - z + z^2)}.$$

*Proof.* It follows from the relation  $t(z) = \frac{1}{zr} + 1$ . □

It seems difficult to obtain a close form for the coefficient  $g(n, k)$  using the Lagrange Inversion Formula. Indeed,  $t(z)$  can be expressed in terms of  $C(u)$ , where  $u = \frac{1}{4}(-z^6 + 2z^4 - 2z^3 + 3z^2 + 2z)$  is a polynomial of degree 6, which complicates the calculations.

## 4.2 A bijective approach

Corollary 11 proves that the set of  $UU$ -less Motzkin paths with air pockets of length  $n$  (ending on the  $x$ -axis) is equinumerous to the set of Motzkin paths of length  $n$  avoiding  $UHU$ . Below, we provide a bijection between these two sets.

Any  $UU$ -less Motzkin path with air pockets is either of the form (i)  $\epsilon$ , (ii)  $H\alpha$ , (iii)  $UD\alpha$ , (iv)  $UHD\alpha$ , (v)  $UH\alpha HD\beta$ , or (vi)  $UH^k\gamma D_i\beta$ , where  $\alpha, \beta$  are  $UU$ -less MAP (possibly empty), and  $\gamma D_{i-1}$  is a  $UU$ -less MAP and  $k \geq 1$ . According to all these cases, we define the map  $\chi$ .

**Definition 3** We recursively define the map  $\chi$  from  $UU$ -less MAP of length  $n$  to  $UHU$ -less Motzkin paths of length  $n$ . Let  $P$  be a  $UU$ -less MAP, we set:

$$\chi(P) = \begin{cases} \epsilon & \text{if } P = \epsilon, & (i) \\ H\chi(\alpha) & \text{if } P = H\alpha, & (ii) \\ UD\chi(\alpha) & \text{if } P = UD\alpha, & (iii) \\ UHD\chi(\alpha) & \text{if } P = UHD\alpha, & (iv) \\ UHH\chi(\alpha)D\chi(\beta) & \text{if } P = UH\alpha HD\beta, & (v) \\ U\chi(\gamma D_{i-1})H^{k-1}D\chi(\beta) & \text{if } P = UH^k\gamma D_i\beta, & (vi) \end{cases}$$

Due to the recursive definition, the image of  $UU$ -less MAP of length  $n$  by  $\chi$  is a  $UHU$ -less Motzkin path of length  $n$ . The recursive definition naturally induces that  $\chi$  is a bijection. For instance, if  $P = UHUHD_2UDUHUHHUD_3H$  then we obtain  $\chi(P) = \chi(UH^1UHD_2)\chi(UD)\chi(UHUHHUD_3)\chi(H) = UUHDDUDUUDHDDH$  (see Figure 6 for an illustration of this example).




Figure 6:  $\chi(P) = UUHDDUDUUDHDDH$ .

## 5 Going further

In [1] and [9] some interesting results on the generating functions of restricted lattice paths are obtained for a finite set of types of moves. Can we adapt these techniques to Motzkin paths with air pockets (avoiding some patterns? Considering [3, 4], is it possible to obtain an efficient algorithm for the exhaustive generation of partial Motzkin paths with air pockets? Is it possible to list PMAP of a given length in Gray code order using [5, 10]?

## 6 Statements and Declarations

**Competing Interests:** Authors have non-financial interests that are directly or indirectly related to the work submitted for publication.

**Funding:** The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

## References

- [1] A. Asinowski, C. Banderier, and V. Roitner. Generating functions for lattice paths with several forbidden patterns. *Sémin. Lothar. Comb.* (2020), 84B.
- [2] C. Banderier, M. Bousquet-Mélou, A. Denise, P. Flajolet, and D. Gouyou-Beauchamps. Generating functions for generating trees. *Discrete Math.* **246** (2002), 29–55.
- [3] E. Barcucci, A. Bernini, and R. Pinzani. Exhaustive generation of some lattice paths and their prefixes. *Theoret. Comput. Sci.* **878-879** (2021), 47–52.
- [4] E. Barcucci, A. Bernini, and R. Pinzani. Exhaustive generation of positive lattice paths. *CEUR Workshop proceedings*, **2113** (2018), 79–86.
- [5] E. Barcucci, A. Bernini, and R. Pinzani. A Gray code for regular languages. *CEUR Workshop proceedings*, **2113** (2018), 87–93.
- [6] J.-L. Baril, S. Kirgizov, R. Maréchal, and V. Vajnovszki. Enumeration of Dyck paths with air pockets. *J. Integer Seq.* **26** (2023), Article 23.3.2.
- [7] J.-L. Baril, S. Kirgizov, R. Maréchal, and V. Vajnovszki. Grand Dyck paths with air pockets. *Art Discrete Appl. Math.* **7** (2024), # P1.07.
- [8] J.-L. Baril and P. Barry. Two kinds of partial Motzkin paths with air pockets. *Ars Math. Contemp.* 2024.
- [9] C. Bean, A. Bernini, M. Cervetti, and L. Ferrari. On the generating functions of pattern-avoiding Motzkin paths. *J. Symbolic Comput.* **113** (2022), 126–138.
- [10] A. Bernini, S. Bilotta, R. Pinzani, and V. Vajnovszki. A trace partitioned Gray code for  $q$ -ary generalized Fibonacci strings. *J. Discret. Math. Sci. Cryptogr.*, **18** (2015), 751–761.
- [11] T.-X. He and R. Sprugnoli. Sequence characterization of Riordan arrays. *Discrete Math.* **309** (2009), 3962–3974.
- [12] Q. Hou and T. Mansour. Kernel method and linear recurrence system. *J. Comput. Appl. Math.*, **261**(1) (2008), 227–242.
- [13] A. Krinik, G. Rubino, D. Marcus, R.J. Swift, H. Kasfy, and H. Lam. Dual processes to solve single server systems. *Journal of Stat. Planning and Inference*, **135**(1) (2005), 121–147.
- [14] S.-C. Liu, J. Ma, and Y.-N. Yeh. Dyck paths with peak- and valley- avoiding sets. *Stud. Appl. Math.* **121** (2008), 263–289.
- [15] T. Mansour, M. Schork, and Y. Sun. Motzkin numbers of higher rank: Generating function and explicit expression, *J. Integer Seq.*, **10** (2007), Article 07.7.4.

- [16] D. Merlini, R. Sprugnoli, and M. C. Verri. Lagrange inversion: when and how, *Acta Appl. Math.* **94** (2006), 233–249.
- [17] T. Mansour and M. Shattuck. Counting humps and peaks in generalized Motzkin paths, *Discrete Appl. Math.*, **161** (2013), 2213–2216.
- [18] D. Merlini, D. G. Rogers, R. Sprugnoli, and M. C. Verri. On some alternative characterizations of Riordan arrays. *Canadian J. Math.* **49** (1997), 301–320.
- [19] D. Merlini, D. G. Rogers, R. Sprugnoli, and M. C. Verri. Underdiagonal lattice paths with unrestricted steps. *Discrete Appl. Math.* **91** (1999), 197–213.
- [20] H. Prodinger. The kernel method: a collection of examples. *Sém. Lothar. Combin.* **50** (2004), B50f.
- [21] H. Prodinger. Partial Dyck paths with air pockets. *Integers* **22** (2022), #A24.
- [22] J. L. Ramírez and V. F. Sirvent. A generalization of the  $k$ -bonacci sequence from Riordan arrays. *Electron. J. Combin.* **22** (2015), # P1.38.
- [23] J. L. Ramírez and V. F. Sirvent. Generalized Schröder matrix and its combinatorial interpretation. *Linear Multilinear Algebra* **66**(2) (2018), 418–433.
- [24] D. G. Rogers. Pascal triangles, Catalan numbers and renewal arrays. *Discrete Math.* **22** (1978), 301–310.
- [25] L. W. Shapiro, S. Getu, W. Woan, and L. Woodson, The Riordan group, *Discrete Appl. Math.* **34** (1991), 229–239.
- [26] L. W. Shapiro, R. Sprugnoli, P. Barry, G.-S. Cheon, T.-X. He, D. Merlini, and W. Wang. *The Riordan Group and Applications*, Springer Monographs in Mathematics, Springer, 2022.
- [27] N. J. A. Sloane. The On-line Encyclopedia of Integer Sequences. Available electronically at <http://oeis.org>.
- [28] H. Wilf. *Generating functionology*, Academic Press, New York, 1990.
- [29] S.-L. Yang, Y.-N. Dong, and T.-X. He. Some matrix identities on colored Motzkin paths. *Discrete Math.* **340** (2017), 3081–3091.
- [30] S.-L. Yang and M. Jiang. The  $m$ -Schröder paths and  $m$ -Schröder numbers. *Discret. Math.* **344**(2) (2021), 112209.
- [31] L. Yang, S.-L. Yang, and T.-X. He. Generalized Schröder matrices arising from enumeration of lattice paths. *Czechoslovak Math. J.* **70** (2020), 411–433.