

FIBONACCI AND CATALAN PATHS IN A WALL

JEAN-LUC BARIL AND JOSÉ L. RAMÍREZ

ABSTRACT. We study the distribution of some statistics defined for paths contained in walls. We present the results by giving generating functions, asymptotic approximations, and as well as some closed combinatorial formulas. We prove algebraically that paths in walls of a given width and ending on the x -axis are enumerated by the Catalan numbers, and we provide a bijection between these paths and Dyck paths. We also find that paths in walls with a given number of steps are enumerated by the Fibonacci numbers. Finally, we give a constructive bijection between the paths in walls of a given length with peakless Motzkin paths of a given length.

1. INTRODUCTION AND NOTATION

Lattice path theory takes an important place in combinatorics. In the literature, there are many articles that study combinatorial problems on lattice paths (see [15]). Most of the time, lattice paths are defined in \mathbb{Z}^2 by a starting point (almost always the origin), and a sequence of vectors (also called steps) lying in a given set S . For instance, paths (also called walks) defined with $S = \{N, S, E, W\}$, where $N = (0, 1)$, $S = (0, -1)$, $E = (1, 0)$, and $W = (-1, 0)$, are widely studied in \mathbb{N}^2 (see [9, 16, 17] for instance). These paths may overlap themselves and the problem of their enumeration is very interesting but also often difficult to solve whenever two boundary constraints are imposed. On the other hand, lattice paths can be considered in \mathbb{N}^2 by banning the overlaps and by forcing the paths to go to the right; this is the subclass of *directed paths*. For example, if $S = \{U, D\}$, where $U = (1, 1)$, $D = (1, -1)$, then the paths in \mathbb{N}^2 starting at the origin and ending on the x -axis are the famous Dyck paths that are counted with respect to the semilength (number of steps divided by 2) by the Catalan numbers $C_n = \frac{1}{n+1} \binom{2n}{n}$ (see A000108 in Sloane's On-line Encyclopedia of Integer Sequences [26]). Moreover, if we permit steps $H = (1, 0)$, then we obtain the class of Motzkin paths that are enumerated by the sequence A001006 in [26]. We refer to [1, 4, 5, 6, 7, 8, 10, 12, 18, 19, 24, 25] for several works on the enumeration and the generation of such paths (with and without overlaps) with respect to the length and various statistics.

In this work we introduce a new class of paths in \mathbb{N}^2 induced by a regular tiling of the first quadrant: the *wall*. More precisely, a *wall* is a tiling of \mathbb{N}^2 using tiles (or bricks) of size 1×2 organized as shown in Figure 1. Notice that this tiling can be also viewed as the cell

Date: November 21, 2023.

2010 Mathematics Subject Classification. 05A15, 05A19.

Key words and phrases. Catalan, Fibonacci, Dyck path, generating function, kernel method.

structure of the plant tissues (see [27] for instance), and our work studies the enumeration of paths of the sap according to several parameters defined below.




FIGURE 1. The wall tiling of \mathbb{N}^2 .

A *path* in a wall is a lattice path in \mathbb{N}^2 starting at the origin $(0, 0)$ where each step links two close corners of the bricks, by following the sides of the bricks with no overlap and no return to the left (each step touching exactly two corners of some bricks at its beginning and its end). So, a path consists of step $N = (0, 1)$, $S = (0, -1)$, and $E \in \{E_1 = (1, 0), E_2 = (2, 0)\}$ and their connections are constrained by the tiling (E_2 is used on the x -axis and E_1 above). Let \mathcal{P} be the set of all paths in a wall. For instance, Figure 2 shows the two paths $NEEENESESENEES$ and $NEEENESESENEENEN$, which can also be written as $NE_1E_1E_1NE_1E_1SE_1SE_2NE_1E_1S$ and $NE_1E_1E_1NE_1E_1SE_1SE_2NE_1NE_1N$. The first path ends on the x -axis and the second path ends at ordinate 3. Notice that some works [13, 14] have been investigated the connection between paths and tilings of the plane, but this does not correspond with our definition of the paths in a wall.

A *statistic* on the set \mathcal{P} is a function \mathbf{w} from \mathcal{P} to \mathbb{N} . Below, we define three important statistics for our study. The *width* of a path P , denoted $\text{width}(P)$, is the abscissa of its last point. For instance Figure 2 shows two paths of width 10. The *length* of a path P , denoted $\text{length}(P)$, is the length of the path considering as a curve in \mathbb{R}^2 . Figure 2 shows two paths of length 16 and 17, respectively. The *number of steps* of a path P , denoted $\text{nbstep}(P)$, is the number of steps in the path (or equivalently the number of connections of two corners). Figure 2 shows two paths with 15 and 16 steps, respectively.




FIGURE 2. Two paths in a wall $NEEENESESENEES$ and $NEEENESESENEENEN$. The left path ends on the x -axis, its width is 10, it has 15 steps, and its length is 16. The right path ends at ordinate 3, its width is 10, it has 16 steps, and its length is 17.

Now, let us assume that the statistic \mathbf{w} returns either the width, or the length, or the number of steps of a path. For $k \geq 0$, we consider the generating function $f_k = f_k(z)$

(resp. $g_k = g_k(z)$, resp. $h_k = h_k(z)$), where the coefficient of z^n in the series expansion is the number of paths $P \in \mathcal{P}$ such that $w(P) = n$, ending at ordinate k with an up-step N (resp., with a down-step S , resp., with a horizontal-step E). Let f_k^0 (resp. f_k^1) be the generating function consisting of the terms z^n in f_k such that $n + k = 0 \pmod 2$ (resp., $n + k = 1 \pmod 2$). Similarly, we define $g_k^0, g_k^1, h_k^0,$ and h_k^1 . Obviously, we have $f_k = f_k^0 + f_k^1$, $g_k = g_k^0 + g_k^1$, and $h_k = h_k^0 + h_k^1$ for any $k \geq 0$.

Also, we introduce the bivariate generating functions for $i \in \{0, 1\}$,

$$F^i(u, z) = \sum_{k \geq 0} u^k f_k^i(z), \quad G^i(u, z) = \sum_{k \geq 0} u^k g_k^i(z), \quad \text{and} \quad H^i(u, z) = \sum_{k \geq 0} u^k h_k^i(z).$$

For short, we use the notation $F^i(u)$, $G^i(u)$, and $H^i(u)$, $i \in \{0, 1\}$, for these functions.

We will use all these notations for the three following sections of this study according to the choice of the statistic w ($w = \text{width}$ in Section 3, $w = \text{nbstep}$ in Section 4, and $w = \text{length}$ in Section 5).

Outline of the paper. In this paper, we investigate the enumeration problem of the paths defined by the wall with respect to several parameters. In Section 2, we count paths of a given width (ending on a given abscissa) according to the type of the last step and the ordinate of the last point. We provide an asymptotic approximation for the expected ordinate of the last point, and we prove that such paths ending on the x -axis are counted by the well known Catalan numbers. We exhibit a bijection between these paths and Dyck paths. Notice that this last result was already found by Odlyzko [21] in the context of the enumeration of fountains with a given number of coins on the basis. *En passant*, Odlyzko also enumerates fountains with n coins, which allows us to say that paths ending on the x -axis in the wall and having a given number of bricks below the path, are counted by the infinite continued fraction

$$\frac{1}{1 - \frac{z}{1 - \frac{z^2}{1 - \frac{z^3}{\ddots}}}}$$

In this section we also enumerate the paths ending on x -axis with a given area and width. We note that the total area of the paths is related with the path length in binary trees.

In Section 3, we count paths of a given number of steps according to the type of the last step and the ordinate of the last point. We prove that such paths are counted by the Fibonacci numbers. We exhibit a bijection between these paths and binary words avoiding two consecutive ones.

Finally, in Section 4, we make an analogous study for paths having a given length. We prove that such paths are counted by the generalized Catalan number, which are known to also count RNA structures. We exhibit a bijection between these paths and peakless Motzkin paths. Notice that the study made in this last section is equivalent to the study of paths of a given length in the honeycomb (i.e. hexagonal) lattice (it suffices to expand all bricks of the wall into hexagonal cells), which is not the case for the previous sections.

2. PATHS OF A GIVEN WIDTH

In this part, we count paths in \mathcal{P} of a given width, i.e., ending at a given abscissa.

By convention, we fix $f_0^0 = 1$ for considering the empty path consisting of the origin $(0, 0)$ only. Observing that a step $N = (0, 1)$ in a path cannot end on a corner (n, k) , $k \geq 1$, with $n + k = 0 \pmod{2}$, we deduce $f_k^0 = 0$ for $k \geq 1$. Of course, a path ending with N cannot end at ordinate 0, which implies $f_0^1 = 0$. A path ending with N at ordinate $k = 1$ ends necessarily at an even abscissa, and then, it equals either N or QN , where Q is a path ending with a horizontal step $(2, 0)$, which implies $f_1^1 = 1 + h_0^0$. Finally, a path ending with N on a corner (n, k) , $k > 1$, $n + k = 1 \pmod{2}$, follows necessarily an horizontal step $(1, 0)$ that ends at $(n, k - 1)$, which implies $f_k^1 = h_{k-1}^0$, $k > 1$.

Other recurrence relations for g_k^i and h_k^i , $k \geq 0$, $i \in \{0, 1\}$, can be obtained *mutatis mutandis*. So, we obtain the following equations:

$$(1) \quad \begin{cases} \left\{ \begin{array}{l} f_0^0 = 1 \text{ and } f_k^0 = 0, \\ f_0^1 = 0, f_1^1 = 1 + h_0^0, \text{ and } f_k^1 = h_{k-1}^0, \end{array} \right. & k \geq 1, \quad \left\{ \begin{array}{l} g_k^0 = h_{k+1}^1, \\ g_k^1 = 0, \end{array} \right. & \begin{array}{l} k \geq 0, \\ k \geq 0, \end{array} \quad \text{and} \\ \left\{ \begin{array}{l} h_0^0 = z^2 + z^2(h_0^0 + g_0^0) \text{ and } h_k^0 = z(h_k^1 + f_k^1), \\ h_0^1 = 0 \text{ and } h_k^1 = z(h_k^0 + g_k^0), \end{array} \right. & \begin{array}{l} k \geq 1, \\ k \geq 1. \end{array} \end{cases}$$

Summing the recursions in (1), we have:

$$\begin{cases} F^0(u) = 1, \\ F^1(u) = u + \sum_{k \geq 1} u^k h_{k-1}^0 = u + uH^0(u), \end{cases} \quad \begin{cases} G^0(u) = \sum_{k \geq 0} u^k h_{k+1}^1 = \frac{1}{u} H^1(u), \\ G^1(u) = 0, \end{cases},$$

$$\begin{cases} H^0(u) = h_0^0 + z \sum_{k \geq 1} u^k (f_k^1 + h_k^1) \\ \quad = h_0^0 + z(F^1(u) + H^1(u)), \\ H^1(u) = z \sum_{k \geq 0} u^k (g_k^0 + h_k^0) - z(h_0^0 + g_0^0) \\ \quad = z(H^0(u) + G^0(u)) - \frac{h_0^0 - z^2}{z}. \end{cases}$$

Solving the above functional equations, we deduce

$$F^0(u) = 1 \quad \text{and} \quad F^1(u) = \frac{u(h_0^0 z - u + z)}{u^2 z - u + z},$$

$$G^1(u) = 0 \quad \text{and} \quad G^0(u) = -\frac{h_0^0 u z + h_0^0 z^2 + z^2 - h_0^0}{(u^2 z - u + z) z},$$

$$H^0(u) = \frac{z(-u^2 + h_0^0)}{u^2 z - u + z} \quad \text{and} \quad H^1(u) = -\frac{u(h_0^0 u z + h_0^0 z^2 + z^2 - h_0^0)}{(u^2 z - u + z) z}.$$

In order to compute h_0^0 , we use the kernel method [2, 3, 23, 24] on $F^1(u)$. This method consists in cancelling the denominator of $F^1(u)$ by finding u as an algebraic function r of z . So, if we substitute u with r in the numerator then it necessarily equals zero (in order to counterbalance the cancellation of the denominator), which induces the value of h_0^0 .

Thus, we factorize the denominator $u^2z - u + z = z(u - r)(u - s)$ with

$$r = \frac{1 - \sqrt{1 - 4z^2}}{2z} \quad \text{and} \quad s = \frac{1 + \sqrt{1 - 4z^2}}{2z}.$$

Notice that r is the generating function of Catalan numbers, which ensures us that we remain in the ring of formal power series.

So, we obtain

$$h_0^0 = \frac{r - z}{z}.$$

Now, substituting h_0^0 with its value in the above generating functions, and simplifying by $(u - r)$ in the numerator and the denominator, we obtain the following:

Theorem 1. *We have*

$$F^0(u) = 1, \quad F^1(u) = \frac{u}{z(s - u)}, \quad G^0(u) = \frac{z - r}{z^2(u - s)}, \quad G^1(u) = 0, \quad \text{and}$$

$$H^0(u) = \frac{r + u}{s - u}, \quad H^1(u) = \frac{u(r - z)}{z^2(s - u)}.$$

Finally, the bivariate generating function $S(z, u)$, where the coefficient of $z^n u^k$ is the number of paths of width n ending at ordinate k , satisfies

$$S(z, u) = \frac{r(1 + u)}{z^2(s - u)}.$$

The first terms of the series expansion of $S(z, u)$ are

$$\begin{aligned} &1 + u + (u^2 + u)z + (u^3 + u^2 + 2u + 2)z^2 + (u^4 + u^3 + 3u^2 + 3u)z^3 + \\ &+ (u^5 + u^4 + 4u^3 + 4u^2 + 5u + 5)z^4 + (u^6 + u^5 + 5u^4 + 5u^3 + 9u^2 + 9u)z^5 + \\ &+ (u^7 + u^6 + 6u^5 + 6u^4 + 14u^3 + 14u^2 + 14u + 14)z^6 + O(z^7). \end{aligned}$$

Corollary 1. *We have*

$$[u^k]F^0(u) = [k = 0], \quad k \geq 0, \quad [u^k]F^1(u) = \frac{r^k}{z}, \quad k \geq 1,$$

$$[u^k]G^0(u) = \frac{(r - z)r^{k+1}}{z^2}, \quad k \geq 0,$$

$$[u^k]H^0(u) = \frac{r^{k+1}}{z} - 1, \quad k \geq 1, \quad [u^k]H^1(u) = \frac{(r - z)r^k}{z^2}, \quad k \geq 0,$$

$$[u^k]S(z, u) = \frac{r^{k+1}(r + 1)}{z^2}, \quad k \geq 0, \quad \text{and}$$

all other coefficients are equal to zero.

In order to provide closed forms for the coefficients of $z^n u^k$ in the previous generating functions, we need to obtain a closed form for the coefficient of z^n in r^k , $k \geq 0$. The quantity r satisfies the functional equation $r = z(1 + r^2) = z\phi(r)$ with $\phi(t) = 1 + t^2$. From Lagrange inversion, (see [20] for instance), we have

$$[z^n]r^k = \frac{k}{n}[t^{n-k}]\phi(t)^n = \frac{k}{n}[t^{n-k}](1 + t^2)^n.$$

So, we have

$$[z^n]r^k = 0 \text{ if } n - k \not\equiv 0 \pmod{2} \quad \text{and} \quad [z^n]r^k = \frac{k}{n} \binom{n}{\frac{n-k}{2}} \text{ otherwise.}$$

Therefore, we obtain the following.

Theorem 2. *The number $f(n, k)$ of paths of width n ending at ordinate k with a step N is given by*

$$f(n, k) = \frac{k}{n+1} \binom{n+1}{\frac{n+1-k}{2}} \quad \text{if } n+k \not\equiv 0 \pmod{2} \text{ and } 0 \text{ otherwise.}$$

The number $g(n, k)$ of paths of width n ending at ordinate k with a step S is given by

$$g(n, k) = \frac{k+2}{n+2} \binom{n+2}{\frac{n-k}{2}} - \frac{k+1}{n+1} \binom{n+1}{\frac{n-k}{2}} \quad \text{if } n+k \equiv 0 \pmod{2} \text{ and } 0 \text{ otherwise.}$$

Theorem 3. *The number $s(n, k)$ of paths of width n ending at ordinate k is given by*

$$\begin{aligned} s(n, k) &= \frac{k+2}{n+2} \binom{n+2}{\frac{n-k}{2}} \quad \text{if } n+k \equiv 0 \pmod{2} \text{ and} \\ s(n, k) &= \frac{k+1}{n+2} \binom{n+2}{\frac{n-k+1}{2}} \quad \text{otherwise.} \end{aligned}$$

From Theorem 2 and Theorem 3, we can easily deduce a closed form for the number $h(n, k)$ of paths of width n ending at ordinate k with a horizontal step. As a byproduct of Theorem 3, if we set $k = 0$ and $n = 2m$ is even, then $s(2m, 0) = \frac{1}{m+1} \binom{2m+2}{m} = \frac{1}{m+2} \binom{2m+2}{m+1}$, which corresponds to the $(m+1)$ -th Catalan number (see A000108). Figure 3 shows the 14 paths of width 6 ending on the x -axis (i.e, ending at ordinate $k = 0$).




FIGURE 3. The 14 paths of width 6 ending on the x -axis (Catalan number).

Corollary 2. *The generating function for the number of paths of a given width is*

$$S(z, 1) = \frac{2r}{z^2(s-1)} = \frac{1 - z - 2z^2 - (1 - z)\sqrt{1 - 4z^2}}{z^3(-1 + 2z)},$$

and the n -th coefficient in the series expansion is given by

$$2 \cdot \binom{n+1}{\lfloor n/2 \rfloor},$$

which corresponds to twice the coefficients of [A037952](#) in [26].

The first terms of the series expansion are

$$2 + 2z + 6z^2 + 8z^3 + 20z^4 + 30z^5 + 70z^6 + 112z^7 + 252z^8 + 420z^9 + O(z^{10}).$$

By calculating $\partial_u(S(z, u))|_{u=1}$, and using classical methods [11, 22] for an asymptotic approximation of the coefficient of z^n , we obtain the following.

Corollary 3. *An asymptotic for the expected ordinate of the last point in all paths of a given width is given by*

$$\sqrt{\frac{\pi n}{2}} \sim 1.253314137\sqrt{n}.$$

We end this section by exhibiting a constructive bijection ϕ between paths of width $2n$ ending on the x -axis in \mathcal{P} and Dyck paths with $n+1$ up steps. Recall that $E_1 = (1, 0)$ and $E_2 = (2, 0)$. Let \mathcal{P}^1 be the set of paths in the wall starting at $(1, 1)$, ending at ordinate 1, and never going to the x -axis. If Q is a path in \mathcal{P}^1 , then we define the path \bar{Q} in \mathcal{P} obtained from Q after a translation of vector $(-1, -1)$ (notice that some occurrences of E_1E_1 in Q can be transformed by the translation into a step E_2). For instance, if $P = NE_1E_1E_1NE_1E_1SE_1S$, then $Q = E_1E_1NE_1E_1S$ and $\bar{Q} = E_2NE_1E_1S$.

Definition 1. We recursively define the map ϕ from \mathcal{P} to the set \mathcal{D} of Dyck paths as follows. For $P \in \mathcal{P}$, we set:

$$\phi(P) = \begin{cases} UD & \text{if } P = \epsilon, \\ UD\phi(Q) & \text{if } P = E_2 Q \text{ with } Q \in \mathcal{P}, \\ U\phi(\bar{Q})D & \text{if } P = NE_1QE_1S \in \mathcal{P}, \text{ and } Q \in \mathcal{P}^1, \\ U\phi(\bar{Q})D\phi(R) & \text{if } P = NE_1QE_1SE_2R \in \mathcal{P}, \text{ with } R \in \mathcal{P} \text{ and } Q \in \mathcal{P}^1. \end{cases}$$

Due to the recursive definition, the image by ϕ of a path ending on the x -axis of width $2n$ in \mathcal{P} is a Dyck path of semilength $n + 1$, and it is easy to see that ϕ is a bijection. For instance, the image of $NE_1E_1E_1NE_1E_1SE_1SE_2NE_1E_1S$ (of width 10) is the Dyck path $U\phi(E_2NE_1E_1S)D\phi(NE_1E_1S) = UUD\phi(NE_1E_1S)D\phi(NE_1E_1S) = UUDUUDDDUUD$ (of length 12), see Figure 4. We also refer to Figure 5 for an illustration of the last three cases used in the definition of ϕ .




FIGURE 4. The path $NEEENESESENEES$ of width 10 and its image by ϕ , the Dyck path $UUDUUDDDUUD$ of length 12,

2.1. The total area. The *area* of a path P ending on the x -axis is defined as the number of bricks between the path and the x -axis. It is denoted by $\text{area}(P)$. For example, the path in Figure 2 has area 5. In this part, we enumerate the paths in \mathcal{P} of a given width, ending on the x -axis, and with a given area.

The following theorem provides a functional equation satisfied by the generating function

$$F(z, q) = \sum_{P \in \mathcal{P}} z^{\text{width}(P)} q^{\text{area}(P)}.$$

Theorem 4. The bivariate generating function $F(z, q)$, where the coefficient of $z^n q^k$ is the number of paths of width n ending on the x -axis and area k , satisfies the functional equation

$$(2) \quad F(z, q) = 1 + z^2 F(z, q) + z^2 q^2 F(zq, q) + z^4 q^2 F(zq, q) F(z, q).$$

Proof. In the bijection introduced in Definition 1 we note that any non-empty path can be decomposed as E_2Q , NE_1QE_1S , or $NE_1QE_1SE_2R$, where $Q \in \mathcal{P}^1$ and $R \in \mathcal{P}$, see Figure 5 for a pictorial representation. From this decomposition follows the functional equation.




FIGURE 5. Decomposition of a path into E_2Q , or NE_1QE_1S , or $NE_1QE_1SE_2R$.

□

Theorem 5. *An expression for the generating function $F(z, q)$ is given by the continued fraction*

$$F(z, q) = \frac{1}{-z^2 + \frac{1}{1 + \frac{1}{-q^2z^2 + \frac{1}{1 + \frac{1}{-q^4z^2 + \frac{1}{-q^4z^2 + \frac{1}{\ddots}}}}}}}}.$$

Proof. From Theorem 4 we have

$$F(z, q) = \frac{1 + z^2q^2F(zq, q)}{1 - z^2 - z^4q^2F(zq, q)} = \frac{1}{-z^2 + \frac{1}{1 + z^2q^2F(zq, q)}}.$$

Iterating this expression yields the desired result. □

The first terms of the continued fraction are as follows:

$$1 + (1 + q^2)z^2 + (1 + 2q^2 + q^4 + q^6)z^4 + (1 + 3q^2 + 3q^4 + 3q^6 + 2q^8 + q^{10} + q^{12})z^6 + (1 + 4q^2 + 6q^4 + 7q^6 + 7q^8 + 5q^{10} + 5q^{12} + 3q^{14} + 2q^{16} + q^{18} + q^{20})z^8 + O(z^{10}).$$

The paths corresponding to the boldface in the expansion are displayed in Figure 3. Notice that by considering $1 + q \cdot F(\sqrt{q}, \sqrt{q})$, we obtain the generating function for the paths without steps on the x -axis having a given number of bricks below the path, and we retrieve the results of Odlyzko [21] that enumerates fountains with n coins with the continued

fraction

$$\frac{1}{1 - \frac{z}{1 - \frac{z^2}{1 - \frac{z^3}{\ddots}}}}.$$

Corollary 4. *The generating function of the total area in all paths in \mathcal{P} of a given width and ending on x -axis is*

$$\frac{1 - 3z^2 - (1 - z^2)\sqrt{1 - 4z^2}}{z^4(1 - 4z^2)}.$$

An asymptotic of the n -th coefficient in the series expansion is

$$(1 + (-1)^n)2^{n+1}.$$

Proof. Let $G := \partial_q(F(z, q))|_{q=1}$. Then, by differentiating (2) with respect to q , we obtain

$$(3) \quad G = z^2G + (2z^2 + 2z^4F(z, 1) + z^4G)F(z, 1) + z^2(1 + z^2F(z, 1))H,$$

where $H = \partial_q(F(qz, q))|_{q=1}$. From Theorem 1 we have

$$F(z, 1) = \frac{1 - 2z^2 - \sqrt{1 - 4z^2}}{2z^4} = \sum_{n \geq 0} C_{n+1}z^{2n},$$

where C_n is the n -th Catalan number. From definition of $F(z, q)$ we have

$$H = \partial_q(F(qz, q))|_{q=1} = \sum_{n \geq 0} 2nC_{n+1}z^{2n} + G.$$

Since we have

$$\sum_{n \geq 0} 2nC_{n+1}z^{2n} = 2 \frac{1 - 3z^2 - (1 - z^2)\sqrt{1 - 4z^2}}{z^4\sqrt{1 - 4z^2}},$$

we can substitute this expression in (3), and solving for G , we obtain the desired result. \square

The first terms of the series expansion are

$$2z^2 + 14z^4 + 74z^6 + 352z^8 + 1588z^{10} + 6946z^{12} + 29786z^{14} + 126008z^{16} + O(z^{18}),$$

which corresponds to [A138156](#) in [26], that also counts the sum of the path lengths of all binary trees with n edges. So, it would be interesting to investigate the link between the area in these paths and the path length in binary trees.

3. PATHS WITH A GIVEN NUMBER OF STEPS

In this part, we count paths in \mathcal{P} with a given number of steps (recall that a step is a move $N = (0, 1)$, $S = (0, -1)$, $E \in \{(1, 0), (2, 0)\}$ connecting two close corners in a wall. With the same similar arguments used in the previous section, we easily obtain the following recurrence relations.

$$(4) \quad \begin{cases} f_0^0 = 1, \text{ and } f_k^0 = 0, & k \geq 1, \\ f_0^1 = 0, f_1^1 = z + zh_0^0, \text{ and } f_k^1 = zh_{k-1}^0, & k \geq 2, \end{cases}, \quad \begin{cases} g_k^0 = zh_{k+1}^1, & k \geq 0, \\ g_k^1 = 0, & k \geq 0, \end{cases} \quad \text{and}$$

$$\begin{cases} h_0^0 = z + z(h_0^0 + g_0^0), \text{ and } h_k^0 = z(h_k^1 + f_k^1), & k \geq 1, \\ h_0^1 = 0, \text{ and } h_k^1 = z(h_k^0 + g_k^0), & k \geq 1. \end{cases}$$

Summing the recursions in (4), we have:

$$\begin{cases} F^0(u) = 1, \\ F^1(u) = zu + z \sum_{k \geq 1} u^k h_{k-1}^0 = zu + zuH^0(u), \end{cases} \quad \begin{cases} G^0(u) = z \sum_{k \geq 0} u^k h_{k+1}^1 = \frac{z}{u} H^1(u), \\ G^1(u) = 0, \end{cases}$$

$$\begin{cases} H^0(u) = h_0^0 + z \sum_{k \geq 1} u^k (f_k^1 + h_k^1) \\ \quad = h_0^0 + z(F^1(u) + H^1(u)), \\ H^1(u) = z \sum_{k \geq 0} u^k (g_k^0 + h_k^0) - z(h_0^0 + g_0^0) \\ \quad = z(H^0(u) + G^0(u)) - h_0^0 + z. \end{cases}$$

Solving the above functional equations, we deduce

$$F^0(u) = 1 \text{ and } F^1(u) = \frac{uz(h_0^0 uz + h_0^0 z^2 - h_0^0 u + z^2 - u)}{-uz^4 + u^2 z^2 + uz^2 + z^2 - u},$$

$$G^1(u) = 0 \text{ and } G^0(u) = -\frac{z(h_0^0 uz^2 + h_0^0 z - h_0^0 + z)}{-uz^4 + u^2 z^2 + uz^2 + z^2 - u},$$

$$H^0(u) = \frac{uz^4 - (u^2 - h_0^0 + u)z^2 + h_0^0 uz - h_0^0 u}{-uz^4 + (u^2 + u + 1)z^2 - u} \text{ and } H^1(u) = -\frac{u(h_0^0 uz^2 + h_0^0 z - h_0^0 + z)}{-uz^4 + u^2 z^2 + uz^2 + z^2 - u}.$$

In order to compute h_0^0 , we use the kernel method on $F^1(u)$ (see Section 2 for more details). We factorize the denominator $-uz^4 + u^2 z^2 + uz^2 + z^2 - u = z^2(u-r)(u-s)$ with

$$r = \frac{1 - z^2 + z^4 - \sqrt{(z^2 + z + 1)(z^2 - z + 1)(z^2 + z - 1)(z^2 - z - 1)}}{2z^2} \text{ and}$$

$$s = \frac{1 - z^2 + z^4 + \sqrt{(z^2 + z + 1)(z^2 - z + 1)(z^2 + z - 1)(z^2 - z - 1)}}{2z^2}.$$

Cancelling the numerator of $F^1(u)$ by substituting u with r , we obtain

$$h_0^0 = \frac{z}{1 - z - rz^2}.$$

The first terms of the series expansion are

$$z + z^2 + z^3 + z^4 + 2z^5 + 3z^6 + 5z^7 + 7z^8 + 11z^9 + O(x^{10}),$$

and this sequence does not appear in [26]. However, the sequence of odd powers corresponds to [A051286](#), and the coefficient $h(2n+1)$ of z^{2n+1} is given by

$$h(2n+1) = \sum_{k=0}^n \binom{n-k}{k}^2.$$

The sequence of even powers corresponds to [A203611](#), and the coefficient $h(2n)$ of z^{2n} is given by

$$h(2n) = \sum_{k=0}^n \binom{k-1}{2k-1-n} \binom{k}{2k-n}.$$

Now, substituting h_0^0 by its value in the previous generating functions, and simplifying by $(u-r)$ in the numerators and denominators, we obtain the following:

Theorem 6. *We have*

$$F^0(u) = 1, \quad F^1(u) = \frac{u(h_0^0 z - h_0^0 - 1)}{z(u-s)}, \quad G^0(u) = \frac{-zh_0^0}{u-s}, \quad G^1(u) = 0, \quad \text{and}$$

$$H^0(u) = \frac{ru + h_0^0}{r(s-u)}, \quad H^1(u) = \frac{-uh_0^0}{u-s}.$$

Finally, the bivariate generating function $S(z, u)$, where the coefficient of $z^n u^k$ is the number of paths with n steps ending at ordinate k , satisfies

$$S(z, u) = 1 + \frac{-ru - zru - h_0^0(ru + z^2 r + z)}{zr(u-s)}.$$

The first terms of the series expansion of $S(z, u)$ are

$$1 + (u+1)z + (1+2u)z^2 + (u^2 + 3u + 1)z^3 + (2u^2 + 4u + 2)z^4 + (u^3 + 4u^2 + 5u + 3)z^5 +$$

$$+ (2u^3 + 6u^2 + 8u + 5)z^6 + (u^4 + 5u^3 + 9u^2 + 12u + 7)z^7 +$$

$$+ (2u^4 + 8u^3 + 14u^2 + 20u + 11)z^8 + O(z^9).$$

Theorem 7. *We have*

$$[u^k]F^0(u) = [k=0], \quad k \geq 0, \quad [u^k]F^1(u) = \frac{1 + h_0^0(1-z)}{z} \cdot r^k, \quad k \geq 1,$$

$$[u^k]G^0(u) = zh_0^0 r^{k+1}, \quad k \geq 0,$$

$$[u^k]H^0(u) = (1 + h_0^0)r^k, \quad k \geq 1, \quad [u^k]H^1(u) = h_0^0 r^k, \quad k \geq 1,$$

$$S(z, 0) = h_0^0/z, \quad [u^k]S(z, u) = \frac{1 + 2h_0^0}{z} \cdot r^k, \quad k \geq 1, \quad \text{and}$$

all other coefficients are equal to zero.

In order to provide a closed form for the coefficient of z^n of all these previous quantities, we need to provide a closed form of $r(n, k) := [z^n]r^k$. We set $r' = r(\sqrt{z})$. Then r' is the generating function for the generalized Catalan numbers, and using the comment of Barry in [26] (see A004148) giving the general term of r'^k , we can easily obtain:

$$\begin{aligned} r(2k, k) &:= [z]^{2k}r^k = 1, \quad k \geq 0 \\ r(2k + 2\ell, k) &:= [z]^{2k+2\ell}r^k = k \sum_{i=\lceil \frac{\ell+1}{2} \rceil}^{\ell} \frac{1}{i} \binom{i}{\ell-i} \binom{i+k-1}{\ell+k-i}, \quad k \geq 1, \ell \geq 1, \\ r(2n + 1, k) &:= [z]^{2n+1}r^k = 0, \quad \text{otherwise.} \end{aligned}$$

Corollary 5. *The number $s(n, k)$ of paths with n steps in \mathcal{P} ending at ordinate k is given by*

$$s(n, k) = r(n + 1, k) + 2 \sum_{i \geq 0}^{n+1} h(i)r(n + 1 - i, k),$$

where $r(n, k)$ and $h(n, k)$ are defined previously.

Corollary 6. *The generating function for the number of paths of a given number of steps in \mathcal{P} is*

$$S(z, 1) = \frac{1 + z}{1 - z - z^2},$$

and the n -th coefficient $s(n)$ in the series expansion is given by a shift of the Fibonacci sequence A000045 in [26], defined by $s(n) = s(n - 1) + s(n - 2)$ anchored with $s(0) = 1$ and $s(1) = 2$.

The first terms of the series expansion are

$$1 + 2z + 3z^2 + 5z^3 + 8z^4 + 13z^5 + 21z^6 + 34z^7 + 55z^8 + 89z^9 + O(z^{10}).$$

Figure 6 shows the eight paths with four steps. Notice that there is a simple bijection ψ between the set of paths in \mathcal{P} with n steps and the set of binary words of length n that do not contain two adjacent ones: reading the path from left to right, we replace each vertical step with 1, and each horizontal step with 0. For instance, the image of the path *ENENESEEN* is 0101001001.

Corollary 7. *The generating function for the number of paths of a given number of steps in \mathcal{P} and ending on the x -axis is*

$$\frac{h_0^0}{z} = \frac{2}{1 - 2z + z^2 - z^4 + \sqrt{1 - 2z^2 - z^4 - 2z^6 + z^8}},$$

and the n -th coefficient of the series expansion is given by $h(n + 1)$ (already defined previously).




FIGURE 6. The eight paths with four steps (Fibonacci).

By calculating $\partial_u(S(z, u))|_{u=1}$ and using classical methods [11, 22] for an asymptotic approximation of the coefficient of z^n , we obtain the following.

Corollary 8. *An asymptotic for the expected ordinate of the last point in all paths in \mathcal{P} of a given number of steps is*

$$\frac{\sqrt{-15 + 7\sqrt{5}} (5 + 3\sqrt{5}) \sqrt{n}}{10\sqrt{\pi}} \sim 0.5335775634\sqrt{n}.$$

4. PATHS WITH A GIVEN LENGTH

In this part, we count paths in \mathcal{P} with a given length. So, we easily obtain the following recurrence relations.

$$(5) \quad \begin{cases} f_0^0 = 1 \text{ and } f_k^0 = 0, & k \geq 1, \\ f_0^1 = 0, f_1^1 = z + zh_0^0, \text{ and } f_k^1 = zh_{k-1}^0, & k \geq 2, \end{cases} \quad \begin{cases} g_k^0 = zh_{k+1}^1, & k \geq 0, \\ g_k^1 = 0, & k \geq 0, \end{cases} \quad \text{and}$$

$$\begin{cases} h_0^0 = z^2 + z^2(h_0^0 + g_0^0) \text{ and } h_k^0 = z(h_k^1 + f_k^1), & k \geq 1, \\ h_0^1 = 0 \text{ and } h_k^1 = z(h_k^0 + g_k^0), & k \geq 1. \end{cases}$$

Summing the recursions in (5), we have:

$$\begin{cases} F^0(u) = 1, \\ F^1(u) = zu + z \sum_{k \geq 1} u^k h_{k-1}^0 = zu + zuH^0(u), \end{cases} \quad \begin{cases} G^0(u) = z \sum_{k \geq 0} u^k h_{k+1}^1 = \frac{z}{u} H^1(u), \\ G^1(u) = 0, \end{cases}$$

$$\begin{cases} H^0(u) = h_0^0 + z \sum_{k \geq 1} u^k (f_k^1 + h_k^1) \\ \quad = h_0^0 + z(F^1(u) + H^1(u)), \\ H^1(u) = z \sum_{k \geq 0} u^k (g_k^0 + h_k^0) - z(h_0^0 + g_0^0) \\ \quad = z(H^0(u) + G^0(u)) - \frac{h_0^0 - z^2}{z}. \end{cases}$$

Solving the above functional equations, we deduce

$$F^0(u) = 1 \text{ and } F^1(u) = \frac{uz(h_0^0 z^2 + z^2 - u)}{-uz^4 + u^2 z^2 + uz^2 + z^2 - u},$$

$$G^1(u) = 0 \text{ and } G^0(u) = -\frac{h_0^0 uz^2 + h_0^0 z^2 + z^2 - h_0^0}{-uz^4 + u^2z^2 + uz^2 + z^2 - u},$$

$$H^0(u) = \frac{z^2(uz^2 - u^2 + h_0^0 - u)}{-uz^4 + u^2z^2 + uz^2 + z^2 - u} \text{ and } H^1(u) = -\frac{u(h_0^0 uz^2 + h_0^0 z^2 + z^2 - h_0^0)}{(-uz^4 + u^2z^2 + uz^2 + z^2 - u)z}.$$

In order to compute h_0^0 , we use the kernel method on $F^1(u)$ (see Section 2 for more details). We factorize the denominator $-uz^4 + u^2z^2 + uz^2 + z^2 - u = z^2(u-r)(u-s)$ where r and s are the same as those defined in Section 3.

Cancelling the numerator of $F^1(u)$ by substituting u with r , we obtain

$$h_0^0 = \frac{r - z^2}{z^2}.$$

Now, substituting h_0^0 by its value in the previous generating functions, and simplifying by $(u-r)$ in the numerators and denominators, we obtain the following:

Theorem 8. *We have*

$$F^0(u) = 1, \quad F^1(u) = \frac{-u}{z(u-s)}, \quad G^0(u) = \frac{z^2 - r}{z^2(u-s)}, \quad G^1(u) = 0, \text{ and}$$

$$H^0(u) = \frac{1}{z^2(s-u)} - 1, \quad H^1(u) = \frac{u(z^2 - r)}{z^3(u-s)}.$$

Finally, the bivariate generating function $S(z, u)$, where the coefficient of $z^n u^k$ is the number of paths of length n ending at ordinate k , satisfies

$$S(z, u) = \frac{z^3 - rz - z - ru}{z^3(u-s)}.$$

The first terms of the series expansion of $S(z, u)$ are

$$1 + uz + (u+1)z^2 + (u^2 + 2u)z^3 + (u^2 + 2u + 2)z^4 + (u^3 + 3u^2 + 3u)z^5 +$$

$$+ (u^3 + 3u^2 + 4u + 4)z^6 + (u^4 + 4u^3 + 6u^2 + 6u)z^7 +$$

$$+ (u^4 + 4u^3 + 7u^2 + 9u + 8)z^8 + O(z^9).$$

Theorem 9. *We have*

$$[u^k]F^0(u) = [k=0], \quad k \geq 0, \quad [u^k]F^1(u) = \frac{r^k}{z}, \quad k \geq 1,$$

$$[u^k]G^0(u) = r^{k+2}/z^2 - r^{k+1}, \quad k \geq 0,$$

$$[u^k]H^0(u) = r^{k+1}/z^2, \quad k \geq 1, \quad [u^k]H^1(u) = r^{k+1}/z^3 - r^k/z, \quad k \geq 1,$$

$$S(z, 0) = \frac{r - z^2}{z^4}, \quad [u^k]S(z, u) = \frac{r^{k+1}}{z^2} \left(1 + r + \frac{1}{z} - z^2\right), \quad k \geq 1, \text{ and}$$

all other coefficients are equal to zero.

Using the closed form of $[z^n]r^k$ obtained in the previous section, we deduce the following.

Corollary 9. *The number $s(n, k)$ of paths of length n in \mathcal{P} ending at ordinate k is given by*

$$s(2, 0) = 1, s(n, 0) = r(n + 4, 1), \quad n \geq 4, \quad \text{and}$$

$$s(n, k) = r(n + 2, k + 1) + r(n + 3, k + 1) - r(n, k + 1) + r(n + 2, k + 2).$$

Corollary 10. *The generating function for the number of paths of a given length in \mathcal{P} is*

$$S(z, 1) = \frac{z^3 - rz - z - r}{z^3(1 - s)} = \frac{-1 + 2z^2 + 2z^5 + z^6 + (1 - z^2)\sqrt{1 - 2z^2 - z^4 - 2z^6 + z^8}}{2z^5(1 - z - z^2)}.$$

The sequences of coefficients of z^{2n+1} corresponds to [A003440](#) and the sequences of coefficients of z^{2n} does appear in [26]

Corollary 11. *The generating function for the number of paths of a given length in \mathcal{P} ending on the x -axis is $\frac{r-z^2}{z^4}$ and the n -th coefficient of the series expansion is given by $r(2n, 0)$, which corresponds to the n -th generalized Catalan number (see [A004148](#)).*

Corollary 12. *An asymptotic for the expected ordinate of the last point in all paths in \mathcal{P} of a given length is*

$$\frac{(3 + \sqrt{5}) \sqrt{-15 + 7\sqrt{5}}}{2\sqrt{\pi n}} \sim 1.193115703 \cdot \frac{1}{\sqrt{n}}.$$

Figure 7 shows the eight paths of length 8 ending on the x -axis.




FIGURE 7. The eight paths of length 8 ending on the x -axis (generalized Catalan)

We end the section by exhibiting a constructive bijection ψ between paths of length $2n$ ending on the x -axis in \mathcal{P} and peakless Motzkin paths with $n + 1$ steps (i.e, Motzkin paths with no occurrence of UD). Recall that $E_1 = (1, 0)$ and $E_2 = (2, 0)$, and that \mathcal{P}^1 is the set of paths in the wall starting at $(1, 1)$, ending at ordinate 1, and never going to the x -axis. If $Q \in \mathcal{P}^1$, then we recall that \bar{Q} is the path in \mathcal{P} obtained from Q after the translation of the vector $(-1, -1)$. These definition are already given before the bijection in Section 2.

Definition 2. We recursively define the map ψ from \mathcal{P} to the set \mathcal{PM} of peakless Motzkin paths as follows. For $P \in \mathcal{P}$, we set:

$$\psi(P) = \begin{cases} H & \text{if } P = \epsilon, \\ H\psi(Q) & \text{if } P = E_2 Q \text{ with } Q \in \mathcal{P}, \\ U\psi(\bar{Q})D & \text{if } P = NE_1Q E_1S \in \mathcal{P}, \text{ and } Q \in \mathcal{P}^1, \\ U\psi(\bar{Q})D\psi(R) & \text{if } P = NE_1Q E_1S E_2R \in \mathcal{P}, \text{ with } R \in \mathcal{P} \text{ and } Q \in \mathcal{P}^1. \end{cases}$$

Due to the recursive definition, the image by ψ of a path ending on the x -axis of length $2n$ in \mathcal{P} is a peakless Motzkin path with $n + 1$ steps, and it is easy to see that ψ is a bijection. For instance, the image of $NE_1E_1E_1NE_1E_1SE_1SE_2NE_1E_1S$ (of length 16) is the peakless Motzkin path $UHUHDDUHD$ (of length 9), see Figure 8. Notice that the definition of ψ is basically the same as in Definition 1 since we use the same decomposition of a path.




FIGURE 8. The path $NE_1E_1E_1NE_1E_1SE_1SE_2NE_1E_1S$ of length 16 and its image by ψ , the peakless Motzkin path $UHUHDDUHD$ of length 9.

5. CONCLUSIONS

This work enumerates the width, the number of steps, the length, and the area in all paths contained in a wall. The main tools are multivariate generating functions and the kernel method. We can extend our results by considering additional statistics, for example the number of *turns* of the path, i.e., the number of occurrences of the subpaths EN , ES , NE , and SE . For example, it is possible to give an expression for the multivariate generating function $S(z, u, w)$, where the coefficient of $z^n u^k w^\ell$ is the number of paths with n steps, ℓ turns, and ending at ordinate k . The expression is too large, however, the first few terms of the Taylor expansion are

$$\begin{aligned} S(z, u, w) = & 1 + (1 + u)z + (1 + 2uw)z^2 + (1 + 2uw + uw^2 + u^2w^2)z^3 \\ & + (1 + 2uw + w^2 + 2uw^2 + 2u^2w^3)z^4 \\ & + (1 + 2uw + 3uw^2 + u^2w^2 + 2w^3 + 2u^2w^3 + u^2w^4 + u^3w^4)z^5 + O(z^6). \end{aligned}$$

Moreover, the bivariate generating function $S(z, 1, w)$, where the coefficient of $z^n w^\ell$ is the number of paths with n steps and ℓ turns is given by

$$S(z, 1, w) = \frac{(w^2 - 2w + 1)z^2 - z - 1}{w^2z^2 + z - 1}.$$

Therefore, the generating function for the total number of turns in all paths of a given width is

$$\frac{2z^2(1+z)}{(1-z-z^2)^2} = 2z^2 + 6z^3 + 14z^4 + 30z^5 + 60z^6 + 116z^7 + 218z^8 + 402z^9 + O(z^{10}),$$

and the n -th coefficient corresponds to twice the convolution of the Fibonacci sequence (see [A023610](#) in [26]), that is,

$$2 \cdot \sum_{i=2}^{n+1} F_i F_{n+1-i}.$$

REFERENCES

- [1] A. Asinowski, C. Banderier, and V. Roitner. Generating functions for lattice paths with several forbidden patterns. *Sémin. Lothar. Comb.*, 84B (2020).
- [2] C. Banderier, M. Bousquet-Mélou, A. Denise, P. Flajolet, and D. Gouyou-Beauchamps. Generating functions for generating trees. *Discrete Math.*, 246 (2002), 29–55.
- [3] C. Banderier, and Ph. Flajolet. Basic Analytic Combinatorics of Directed Lattice Paths. *Theoretical Computer Science*, 281(1-2) (2002), 37–80.
- [4] C. Banderier and M. Wallner. Lattice paths with catastrophes. *Discrete Math Theor. Comput. Sci.*, 19(1) (2017), #23.
- [5] E. Barucci, A. Del Lungo, S. Fezzi, and R. Pinzani. Nondecreasing Dyck paths and q -Fibonacci numbers. *Discrete Math.*, 170 (1997), 211–217.
- [6] E. Barucci, A. Bernini, and R. Pinzani. Exhaustive generation of some lattice paths and their prefixes. *Theoret. Comput. Sci.*, 478 (2021), 47–52.
- [7] J.-L. Baril and A. Petrossian. Equivalence classes of Dyck paths modulo some statistics. *Discrete Math.*, 338 (2015), 655–660.
- [8] C. Bean, A. Bernini, M. Cervetti, and L. Ferrari. On the generating functions of pattern-avoiding Motzkin paths. *J. Symbolic Comput.*, 113 (2022), 126–138.
- [9] M. Bousquet-Mélou and M. Mishna. Walks with small steps in the quarter plane. In *Algorithmic probability and combinatorics, vol. 520 of Contemp. Math.*, (2010), Amer. Math. Soc., 1–39.
- [10] E. Deutsch. Dyck path enumeration. *Discrete Math.*, 204 (1999), 167–202.
- [11] P. Flajolet and R. Sedgewick. *Analytic Combinatorics*, Cambridge University Press, 2009.
- [12] R. Flórez and J.L. Ramírez. Enumerating symmetric and asymmetric peaks in Dyck paths. *Discrete Math.*, 343 (2020), 112118.
- [13] S. Heubach and T. Mansour. Staircase tilings and lattice paths. *Congr. Numer.*, 182 (2006), 97–109.
- [14] S. Heubach, N.Y. Li, and T. Mansour. Staircase tilings and k -Catalan structures. *Discrete Math.*, 308(24) (2008), 5954–5953.
- [15] K. Humphreys. A history and a survey of lattice path enumeration. *J. of Stat. Plan. and Inference*, 140(8) (2010), 2237–2254.
- [16] M. Kauers and R. Yatchak. Walks in the quarter plane with multiple steps. In *Proceedings of FPSAC 2015, Discrete Math. Theor. Comput. Sci.*, (2015), 25–36.
- [17] I. Kurkova and K. Raschel. On the functions counting walks with small steps in the quarter plane. *Publ. Math. Inst. Hautes Études Sci.*, 116 (2012), 69–114.
- [18] T. Mansour. Statistics on Dyck paths. *J. Integer Sequences*, 9 (2006), 06.1.5.
- [19] D. Merlini, R. Sprugnoli, and M.C. Verri. Some statistics on Dyck paths, *J. Statis. Plann. Inference*, 101 (2002), 211–227.
- [20] D. Merlini, R. Sprugnoli, and M.C. Verri. Lagrange inversion: when and how. *Acta Appl. Math.*, 94 (2006), 233–249.

- [21] A.M. Odlyzko and H.S. Wilf. The editor's corner: n coins in a fountain. *Amer. Math. Monthly*, 95 (1988), 840–843.
- [22] A.G. Orlov. On asymptotic behavior of the Taylor coefficients of algebraic functions. *Sib. Math. J.*, 25(5) (1994), 1002–1013.
- [23] H. Prodinger. The kernel method: a collection of examples. *Sém. Lothar. Combin.*, 50 (2004), Paper B50f.
- [24] H. Prodinger. Partial skew Dyck paths: a kernel method approach. *Graphs Combin.*, 38 (2022), Paper 135.
- [25] A. Sapounakis, I. Tasoulas, and P. Tsikouras. Counting strings in Dyck paths. *Discrete Math.*, 307 (2007), 2909–2924.
- [26] N. J. A. Sloane. The On-line Encyclopedia of Integer Sequences, available electronically at <http://oeis.org>.
- [27] L.A. Staehelin. Cell wall. *Encyclopedia Britannica*, <https://www.britannica.com/science/cell-wall-plant-anatomy>.

LIB, UNIVERSITÉ DE BOURGOGNE FRANCHE-COMTÉ, B.P. 47 870, 21078, DIJON CEDEX, FRANCE
Email address: `barjl@u-bourgogne.fr`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL DE COLOMBIA, BOGOTÁ, COLOMBIA
Email address: `jlr Ramirezr@unal.edu.co`